

THE EISENHOWER LIBRARY

3 1151 02678 0993

STORAGE

144,682

Library



Johns Hopkins at the University



ON THE REACTIONS OF BOTH THE IONS AND THE
NONIONIZED FORMS OF ELECTROLYTES:
ON THE REACTIONS OF METHYL IODIDE WITH SODIUM,
POTASSIUM AND LITHIUM ETHYLATES AT 0° AND 25°.

DISSERTATION

Submitted to the Board of University Studies of
the Johns Hopkins University in conformity
with the requirements for the degree of
Doctor of Philosophy

by

BESSIE MARION BROWN

Baltimore

June 1913.

THE NEW YORK PUBLIC LIBRARY

ASTOR LENOX TILDEN FOUNDATION

500 MADISON AVENUE, NEW YORK 17, N. Y.

FORM 100 (10-1-60) (REVISED 10-1-60)

166,682

TO: THE NEW YORK PUBLIC LIBRARY
ASTOR LENOX TILDEN FOUNDATION
500 MADISON AVENUE, NEW YORK 17, N. Y.
FROM: THE NEW YORK PUBLIC LIBRARY
ASTOR LENOX TILDEN FOUNDATION
500 MADISON AVENUE, NEW YORK 17, N. Y.

NEW YORK, N. Y.

RECEIVED

1960

CONTENTS.

	Page
Acknowledgment.....	1
Theoretical	
Historical.....	3
Theory of the reactions.....	3
Discussion of the experimental results.....	5
Salt catalysis.....	8
Experimental	
Preparation of methyl iodide.....	15
Analysis of methyl iodide.....	18
Preparation of solutions for salt catalysis.....	20
Calibration of machine.....	20
Determination of loss of alkyl halide.....	23
Technique at 0°.....	34
Tables of reaction velocities.....	37
Conclusions.....	63
Biography.....	65

ACKNOWLEDGMENT

The author takes this opportunity to express her gratitude to Professor Remsen, Professor Moree, Professor Jones, Professor Whitehead, Associate Professor Acree, Associate Professor Lovelace and Dr. Frazer for valuable instruction.

Special thanks are due to Associate Professor Acree at whose suggestion and under whose guidance this work was carried out.

CONFIDENTIAL

The above is a true and correct copy of the original document as submitted to the Department of Defense, Washington, D.C., on the 15th day of May, 1964, by the Department of Defense, Washington, D.C., and is being furnished to you for your information.

Very truly yours,

Special Agent in Charge, Federal Bureau of Investigation, Washington, D.C.

Enclosed are:

ON THE REACTIONS OF BOTH THE IONS AND THE
NONIONIZED FORMS OF ELECTROLYTES:

ON THE REACTIONS OF METHYL IODIDE WITH SODIUM,
POTASSIUM AND LITHIUM ETHYLATES AT 0° AND 35°.

An entirely new theory concerning chemical reaction was brought out by Acree and Brunel¹ in 1905-06, and Acree and Johnson² in 1907, when they showed by experimental evidence that, in some catalytic reactions, the acceleration may take place through the action of both the ionized and nonionized portion of the salt.

Shadinger³, and later, Nirdlinger, Rogers and Chandler, showed that in absolute alcohol both the 1-phenyl-3-thiourazole anions and the nonionized sodium 1-phenyl-3-thiourazole react with alkyl halides.

In working with the phenolates, Robertson⁴ found that both the phenolate ion and the nonionized sodium phenolate react with the alkyl halides, and Marshall, Harrison⁵, and Myers observed that both the ions and molecules of sodium ethylate are concerned in the catalytic effect of this compound on the reversible addition of alcohol to acetonitrile, benzonitrile, para brom benzonitrile and others.

Other reactions studied, and which uphold this theory, are those of methyl iodide, ethyl iodide, and ethyl bromide with sodium, lithium, and potassium ethylates at 35°.

-
1. Am. Chem. J., 43, 505.
 2. Ibid., 38, 359, 360.
 3. Ibid., 39, 375.
 4. Diss., J.H.U., 1910.
 5. Diss., J.H.U., 1912.

OF THE REACTION OF THE ALKYL ALCOHOLS
 WITH THE ALKYL ALCOHOLS
 OF THE ALKYL ALCOHOLS
 OF THE ALKYL ALCOHOLS

An entirely new theory concerning chemical reaction
 was brought out by Jones and Brune¹ in 1905-06, and Jones
 and Johnson² 1907, when they showed by experimental evidence
 that, in some catalytic reactions, the catalysis may take
 place through the action of both the ionized and non-ionized
 portion of the acid.

William³, and later, Winkler⁴, Rogers and Brune⁵,
 showed that in reactions alcohol both the 1-phenyl-2-propanol-
 sole anions and the non-ionized sodium 1-phenyl-2-propanol-
 react with alkyl halides.

In working with the phenolates, Winkler⁴ found that
 both the phenolates and the non-ionized sodium phenolates
 react with the alkyl halides, and Winkler⁴, Johnson⁶, and
 Jones observed that both the ion and non-ionized sodium
 phenolates are concerned in the catalytic effect of this com-
 pound on the reversible reaction of alcohol to acetone,
 ethyl acetate, and other.

Other reactions studied, and which showed that alkyl
 are those of methyl iodide, ethyl iodide, and ethyl bromide
 with sodium, lithium, and potassium phenolates at 50°.

1. J. Jones, J. Brune, J. Am. Chem. Soc., 27, 1005 (1905).
2. J. Jones, J. Johnson, J. Am. Chem. Soc., 29, 1005 (1907).
3. W. Winkler, J. Am. Chem. Soc., 29, 1005 (1907).
4. W. Winkler, J. Am. Chem. Soc., 29, 1005 (1907).
5. J. Rogers, J. Brune, J. Am. Chem. Soc., 29, 1005 (1907).
6. J. Johnson, J. Jones, J. Am. Chem. Soc., 29, 1005 (1907).

Any results which shed light upon the mechanism of organic reactions are of such extreme importance, that it was considered advisable to extend this study to other reactions, and to study some of the same reactions at different temperatures.

With this idea in view, the author took up the study of the reactions of sodium, lithium and potassium ethylates with methyl iodide at 0°, and also the effect of added sodium iodide on the reaction between sodium ethylate and methyl iodide at 35°.

If both the ethylate ion and the nonionized ethylate salt react with the molecular alkyl halide, we have two reactions proceeding simultaneously, one proportional to the ionized part of the ethylate, α , and the other proportional to the nonionized part, $1 - \alpha$. It follows that

$$(1) \quad \frac{dx}{dt} = K_1\alpha(C_{\text{ethylate}} - x)(C_{\text{alkyl halide}} - x) + K_m(1 - \alpha)(C_{\text{ethylate}} - x)(C_{\text{alkyl halide}} - x)$$

in which K_1 is the velocity constant for the ionic reaction, and K_m is the constant for the molecular reaction, in a normal solution, and x is the change in the concentration in the time t . If A represents C_{ethylate} , and B is substituted for $C_{\text{alkyl halide}}$, we obtain by integration the equation

$$(3) \quad K_n = K_1\alpha + K_m(1 - \alpha) = \frac{1}{At} \frac{x}{A - x}, \text{ if } C_{\text{ethylate}} \text{ is equal}$$

to $C_{\text{alkyl halide}}$, or

$$(3) \quad K_n = K_1\alpha + K_m(1 - \alpha) = \frac{1}{(B - A)t} \ln \frac{A(B - x)}{B(A - x)}, \text{ if}$$

C_{ethylate} is not equal to $C_{\text{alkyl halide}}$.

Substituting the data obtained for t , x , A , and B , we ob-

tain a set of constants for K_n for each concentration of ethylate used. K_n represents the molecular velocity constant, or the velocity which that reaction would have if the concentration were normal, but the ionization corresponded to the concentration $1/V$, when V is the number of liters containing one gram molecule. For a complete mathematical and theoretical discussion of these equations, see Johnson and Acree¹, and dissertations of Robertson, and Harrison.

To determine K_i and K_m for the separate ionic and molecular reactions, it is only necessary to substitute the proper values in the following simultaneous equations and solve for K_i and K_m .

$$K_n = K_i\alpha + K_m(1 - \alpha)$$
$$K_n' = K_i\alpha' + K_m(1 - \alpha') \quad \text{etc.}$$

Values for K_i and K_m are thereby obtained for every pair of concentrations, and the average of these is taken as the accepted value.

Robertson discussed the possibility and probability of other reactions taking place simultaneously with those given above. We must consider, for example, the catalytic effect of the salt formed, or even of the ethylate itself, which we could designate as a "salt effect", and perhaps a side reaction, as well, in which olefines may be formed.

Since we obtain, however, from the equations used, two constant values for K_i and K_m , we can state that we have

1. Am. Chem. J., 37, 410; 38, 356; 48, 352.

either two reactions, or two or more sets of reactions, taking place, one being a function of the concentration of the ions, and the other of the nonionized portion.

If K_1 represents the velocity constant of the reaction between the ethylate ion $C_2H_5O^-$ and methyl iodide, we would expect to find experimentally that the value for K_1 is the same whether the methyl iodide is allowed to react with sodium, lithium, or potassium ethylates, although the values for K_m would not necessarily be the same, and might be entirely different.

Probably the best evidence in favor of this theory is the fact that identical values are obtained for the ethylate ion irrespective of the ionization of the ethylate, the character of the metal, and of the value for K_m . For methyl iodide and sodium ethylate at 25° , Robertson obtained the values $K_1 = 0.127$, and $K_m = 0.0594$. The author repeated this work at the very beginning of this investigation, and obtained the values $K_1 = 0.136$ and $K_m = 0.0586$. For potassium ethylate and methyl iodide at 25° , Harrison obtained the values $K_1 = 0.127$ and $K_m = 0.066$. Similar agreement is found for the phenolate ion when potassium, lithium, and sodium phenolates react with methyl iodide and ethyl iodide.

At 0° the velocity of the reaction is very much less, but we should expect to find that the value for K_1 at this temperature is the same whether sodium, lithium, or potassium ethylate is used. Considering the greater difficulty in carrying out these reactions at 0° and the possibility of

values for ϵ_{eff} and ϵ_{eff} are not very different, and the values for ϵ_{eff} are not very different from the values for ϵ_{eff} in the case of the uniaxial crystal.

It is interesting to note that the values for ϵ_{eff} and ϵ_{eff} are not very different from the values for ϵ_{eff} in the case of the uniaxial crystal. This is due to the fact that the values for ϵ_{eff} and ϵ_{eff} are not very different from the values for ϵ_{eff} in the case of the uniaxial crystal. This is due to the fact that the values for ϵ_{eff} and ϵ_{eff} are not very different from the values for ϵ_{eff} in the case of the uniaxial crystal.

Generally the most reliable values of ϵ_{eff} are obtained for the uniaxial crystal. The values for ϵ_{eff} and ϵ_{eff} are not very different from the values for ϵ_{eff} in the case of the uniaxial crystal. This is due to the fact that the values for ϵ_{eff} and ϵ_{eff} are not very different from the values for ϵ_{eff} in the case of the uniaxial crystal. This is due to the fact that the values for ϵ_{eff} and ϵ_{eff} are not very different from the values for ϵ_{eff} in the case of the uniaxial crystal.

greater errors, the agreement between the values of K_1 for the different ethylates is quite satisfactory. Since viscosity may play an important part in the reaction velocities at 0° , especially in the more concentrated solutions, it will be extremely interesting, as well as important, to obtain all the necessary data, so that corrections can be made for the viscosity.

In Tables I-XV of the lithium series the starred values for K_n are obtained by discarding those constants which correspond to reactions which were allowed to run over night. In these cases the reactions were more nearly complete, the result being that the constants had dropped several per cent. A slight drop was always noticed, but this was usually negligible, or very small, in the dilute solutions. For a complete discussion of the probable causes for this drop in the constants during any specific reaction, see the dissertation of Robertson.

To make the constants from the different concentrations of ethylate comparable, the reactions were allowed to run to about the same percentage completion, which was usually 50 to 60 per cent., and an average value for the reaction velocity was obtained by taking the mean of the constants for the different time periods. It will be seen from the results that it makes very little difference in the calculated values for K_n when these starred values are used in place of the other values.

In the case of lithium ethylate the reaction was so much slower than in the other cases, that it was difficult to ob-

tain the desired time periods, since it was of course impossible to titrate by artificial light. For this reason, in some of the more dilute solutions the reactions were 70 to 80 per cent. completed. But since the constants in these solutions did not decrease appreciably, there is practically no error introduced by using these values for the longer time periods.

It can be seen from the tables that there is very little difference between the values of K_n for the different concentrations of potassium ethylate, and that in some of the more dilute solutions, the values vary to such an extent that they nearly overlap for the different concentrations. The only remedy for this is to diminish the error as much as possible by taking the average of a large number of determinations. It was impossible to use solutions more concentrated than N/4 because the potassium iodide formed precipitated out almost immediately, and the ionization of the remaining potassium ethylate consequently changed. It was therefore thought advisable to find values of K_n for 3N/16 and 3N/32 potassium ethylate so as to give a few more calculated values for K_1 and K_m and reduce the error.

By substituting in the equation $K_n = K_1\alpha + K_m(1 - \alpha)$, these values found for K_1 and K_m and the values for α and $1 - \alpha$ corresponding to the different concentrations of the various ethylates, we obtain the theoretical values for K_n given on page 45 under the heading " K_n calculated". These are compared with the corresponding values of " K_n found", and the experimental errors are thus obtained.

It can be seen from the data given that the work at 0° is not quite as satisfactory as that at 25°.

SALT EFFECT.

Although many theories have been advanced concerning the so-called "salt effect" of neutral salts on certain organic reactions, through lack of sufficient data no very satisfactory explanation has yet been reached.

From the results obtained during the last ten years in this laboratory, however, we can at least point out many errors in the earlier work, and present our own conclusions, which seem to be capable of wider application.

The view adopted by Euler¹ in 1900 was in accordance with the belief of Arrhenius² that the salt cation produces this salt effect by causing a change in the ionization of the water. This was the explanation given by Stieglitz to explain the phenomenon of salt catalysis which he found in studying the hydrolysis of imidoester salts³.

In a full discussion of the subject, Acree⁴ has shown that the explanation offered by Euler and Stieglitz is contrary to facts, one of the strongest points against this theory being the experimental data showing that this view calls for a change in the ionization of water many times greater than could possibly be brought about in this way. According to their theory, the same salt in the same concentration should

1. Z. physik. Chem., 32, 348.

2. Ibid., 1, 110; 4, 236; 31, 197.

3. Am. Chem. J., 39, 39, 166, 403, 437, 586, 719. J. Am. Chem. Soc., 32, 331.

4. Am. Chem. J., 41, 475.

500000 22.3

Although many libraries have been destroyed or damaged, the National Library of Medicine has been spared. The National Library of Medicine is the only library in the world that has been spared.

[illegible]

sometimes cause an increase, and sometimes a decrease in the ionization constant of water.

It is believed by Acree, and this idea is shared by others (see Lapworth¹), that salt catalysis is a very complex phenomenon, and involves a large number of factors, both chemical and physical, one of the least important being perhaps the change in ionization of the solvent.

In the same article mentioned above³, some of these different factors which are probably concerned in any salt effect are fully discussed.

In this laboratory, different investigators have studied the effect of certain added salts on the reactions of the alkyl halides with phenolates, ethylates, and urazole salts. They found that the greater part of the effect was perfectly normal in that it corresponded to the change in ionization of the reacting substances, produced by the added salt.

In some cases, however, a small deviation from the calculated amount was observed which was designated as an "abnormal salt effect", and which might be either "positive" or "negative".

Harrison³, in studying the formation of nitriles from imidoesters by using sodium ethylate as a catalyzer, found that the addition of even several moles of sodium iodide lowered the percentage of ionization of the sodium ethylate, and hence the reaction velocity as calculated, which fact shows that the

1. Chem. News, 97, 310.

2. Am. Chem. J., 41, 478-483.

3. Dissertation, J.H.U. 1913.

sometimes occur at intervals, and sometimes a number in the
location of the effect.

It is claimed by some, that this is caused by
others (see page 10), that this occurs in a very small
area, however, and involves a large number of factors,
both chemical and physical, one of the least important being
perhaps the change in location of the effect.

In the case of the mentioned above, some of these
different factors which are probably concerned in any such
effect are fully discussed.

In the laboratory, different in that it is not a
the effect of certain kinds of the effect on the location of the effect,
the effect with chemical, physical, and chemical effects.
The effect of the effect of the effect on the location of the effect,
the effect of the effect of the effect on the location of the effect,
the effect of the effect of the effect on the location of the effect.

In some cases, however, in which the effect of the effect
on the effect of the effect of the effect on the location of the effect,
the effect of the effect of the effect on the location of the effect,
the effect of the effect of the effect on the location of the effect,
the effect of the effect of the effect on the location of the effect.

Therefore, in studying the location of the effect of the effect
of the effect of the effect of the effect on the location of the effect,
the effect of the effect of the effect on the location of the effect,
the effect of the effect of the effect on the location of the effect,
the effect of the effect of the effect on the location of the effect.

"salt effect" here was "normal". Since in this case K_1 and K_m are so nearly equal, even the "normal salt effect" was extremely small.

Robertson¹ found in the reaction of sodium phenolate with certain alkyl halides, that added sodium iodide produced a "positive abnormal salt effect", in other words, the velocity constant was slightly larger than the calculated value.

On the other hand, Chandler³ found that sodium iodide produces practically the calculated effect on the reaction between 1-phenyl-3-thiourazole and ethyl iodide.

The results obtained by adding different amounts of sodium iodide to the reaction between methyl iodide and sodium ethylate at 35° are shown in tables LXXXIII-CXII and in tabulated form on pages 60-2. In the first column will be found the volume containing a gram equivalent of the sodium ethylate. In the second column is the reciprocal of the concentration of the sodium iodide added, and the third column gives the reciprocal of that concentration of sodium ethylate or sodium iodide having the ionization calculated for either of these electrolytes in the mixture by the use of the isohydric principle.

To give a concrete example of the method used in obtaining the calculated values of K_N , let us consider the case of a solution which is N/33 with reference to both sodium ethylate and sodium iodide. According to the isohydric principle, each electrolyte occupies that portion of the solution which

1. Dissertation, J.H.U., 1910.

3. Dissertation, J.H.U., 1913.

"self effect" that was "small". Since in this case γ was
 1, and so nearly equal, even the "small self effect" was not
 really small.

Subsequent work in the region of sodium hydroxide
 with catalysts of high activity, that would require higher pressures
 a "relative secondary self effect", in other words, the velocity
 constant was slightly less than the calculated value.
 On the other hand, generally found that sodium hydroxide
 produced practically the calculated effect on the reaction
 between 1-phenyl-2-thiocarbonyl and vinyl iodide.

The results obtained by adding different amounts of sodium
 iodide to the reaction between vinyl iodide and sodium acrylate
 are shown in Table I. The first column will be found the
 rate of reaction. In the second column is the calculated
 value obtained by using a rate constant of the sodium acrylate
 in the second column is the calculated value of the concentration
 of the sodium iodide added, and the third column gives the
 calculated value of the concentration of sodium iodide to be added
 which having the reaction calculated for either of these
 electrolytes in the mixture of the use of the logarithmic
 value.

To give a concrete example of the method used in obtaining
 the calculated values of k_2 , let us consider the case of
 a solution which is 0.05M with reference to both sodium acrylate
 and sodium iodide. According to the logarithmic relationship
 each electrolyte contributes that portion of the reaction which

is in proportion to the number of ions it furnishes. In this case, then, since the sodium ethylate and sodium iodide are ionized to approximately the same extent, we can say that each of these ionized constituents occupies half of the solution, and the sodium ethylate consequently would have the ionization of a N/16 solution. Then in the equation $K_n = K_1\alpha + K_m(1 - \alpha)$ we would substitute the values of α and $1 - \alpha$ for a N/16 ethylate solution. If the effect of the sodium iodide is perfectly normal, this is the value of K_n which we should obtain experimentally.

It can be seen from the values for K_n found that in every case, the constant obtained is lower than that calculated, the difference being the "auto" or "abnormal" salt effect, which is expressed in the last column/as a percentage of the theoretical constant. ^{on page 61} Since the constant is lower than the theory, the catalytic effect of the salt is negative and must decrease the velocity of the reaction.

With any one concentration of ethylate, as for instance with the N/32 solution, the "auto" effect increases with the concentration of the added iodide, but not in a direct proportion. That is, although the addition of 3/32 mole of sodium iodide produces approximately three times the effect of 1/32 mole of added salt, and 7/32 mole about seven times as great an effect as the 1/32 mole, this relationship does not continue to hold, for 15/32 mole produces only eight times, and 31/32 mole only twelve times as great an effect as does the 1/32 mole of sodium iodide. In similar work with the N/16 sodium ethylate, while 3/16 mole of added salt produces nearly

three times the effect of $1/16$ mole, $7/16$ mole gives only three and one half times, and $15/16$ mole only six times as great an effect.

These figures naturally suggest that the "auto" effect is a function of the ionized portion of the added salt, since the effect decreases with increased concentration in the same manner as would be expected from the suppression of the ionization. It will be shown in equations developed below that the greater portion of the "abnormal" effect of the salt does appear to be a function of the iodide ions.

It must be constantly be borne in mind, in applying this data to such generalizations, that the "abnormal" salt effect is only a very small part of the whole effect of the salt on the reaction velocity, and since the numerical value of this "auto" effect is obtained by subtracting the K_n found from K_n calculated, an error of one per cent. in the velocity constant would constitute an error of fifty per cent. in the "abnormal" salt effect obtained by adding $1/32$ mole of sodium iodide to the $N/32$ sodium ethylate. Consequently from these figures only an approximation can be made concerning the actual effect of the salt, but we can at least obtain some idea of the direction and magnitude of this effect.

By taking the sum of all the percentage lowerings found in the case of $N/32$ ethylate and dividing this sum by 57, since $57/32$ moles of sodium iodide were added in all, we obtain as a result 1.15 which can be considered the average effect of $1/32$ mole of sodium iodide in a $N/32$ ethylate solution. In the same way we obtain 0.86 as the average effect of $1/32$ mole of sodium

These lines the extent of 1/10 mole, 7/10 mole, 1/10 mole, and 1/10 mole only the lines are
these and one half times, and 1/10 mole only the lines are
great an effect.

These figures suggest that the "auto" effect
is a function of the ionization of the added salt, since
the effect decreases with increased concentration in the same
manner as would be expected from the ionization of the ion-
ization. It will be shown in equations derived from the
the "auto" effect of the "auto" effect of the salt more
likely to be a function of the ionic ions.

It must be constantly be borne in mind in applying these
data to such generalizations, that the "auto" effect of the salt is
is only a very small part of the total effect of the salt on
the reaction velocity, and since the numerical value of this
"auto" effect is determined by subtracting the μ from the μ
calculated, an error of one per cent, in the velocity constant
will calculate an error of fifty per cent, in the "auto"
auto effect obtained by adding 1/10 mole of sodium chloride to
the 1/10 sodium chloride. Consequently these values are only
an approximation and are based on assuming the actual effect of
the salt, but we can at least obtain some idea of the direction
of magnitude of the effect.

By taking the sum of all the percentage ionization found
in the case of 1/10 sodium and adding this sum by 1/10, since
1/10 mole of sodium chloride were added in all, we obtain as a
result 1.12 which can be compared to the average effect of 1/10
mole of sodium chloride in a 1/10 sodium chloride. In the case
it would be about 1.02 as the average effect of 1/10 mole of sodium

iodide in the N/16 ethylate, 0.74 for the N/8 ethylate, and 0.65 for the N/4 solutions. It would appear that the salt effect is not merely a function of the added iodide but of the ethylate concentration as well. These facts seem to contradict the Spohr¹ hypothesis. His work seemed to show that a definite concentration of sodium chloride added to different concentrations of hydrochloric acid produced a specific effect on the reaction velocity of the inversion of cane sugar, irrespective of the concentration of the acid, and therefore of the actual velocity of the reaction.

If the $C_2H_5\bar{O}$ and the iodine ion, $\bar{I}$, each have an independent catalytic effect on the two reactions taking place which we represented by K_1 and K_m , we could then express this condition by the following equation,

$$K'_n = [K_1\alpha + K_m(1 - \alpha)][1 + K_sC_s\alpha' + K_eC_e\alpha]$$

in which K'_n represents the actual velocity constant obtained when sodium iodide is added, and K_s and K_e are the constants referring to the catalytic effect of the iodide and ethylate ions respectively. C_s is the concentration of the added salt, α' its ionization, C_e the concentration of the ethylate, and α its ionization. By substituting the found values for K_n and the known values for K_1 , K_m , α , $1 - \alpha$, and α' , we obtain a series of simultaneous equations which can be solved for K_s and K_e . The values for K_s and K_e obtained were constant in most cases. It must be understood that the error in certain cases might be such as to give values for K_s and K_e which did not seem to fit into this scheme. If we take $K_s = -70$ and

1. J. prakt. Chem., 32, 33; 33, 365.

located in the $\frac{1}{2}$ mole, 0.75 for the $\frac{1}{2}$ mole, and 0.50 for the $\frac{1}{2}$ mole. It would appear that the effect of the addition of the added iodide ion to the ethylate concentration as well. These facts seem to confirm the "ion-pair" hypothesis. The work seemed to show that a definite concentration of sodium chloride added to the ethylate concentration of hydrochloric acid produced a marked effect on the reaction velocity of the reaction of some extent, irrespective of the concentration of the acid, and therefore of the actual velocity of the reaction.

If the $\frac{1}{2}$ mole and the iodine ion, $\frac{1}{2}$ mole, are added, the catalytic effect on the two reactions being shown which is represented by V_1 and V_2 , we could then express this condition by the following equation,

$$V_1 = [V_1 + K_1(1 - \alpha)](1 + K_2C_2 + K_3C_3)$$

in which V_1 represents the actual velocity constant, K_1 is the constant of the reaction, K_2 and K_3 are the constants referring to the catalytic effect of the iodide and ethylate ions respectively. C_2 is the concentration of the acid, and C_3 is the concentration of the ethylate, and α is the ionization. By substituting the known values for V_1 and the known values for V_2 , K_1 , α , $1 - \alpha$, and C_2 , we obtain a series of simultaneous equations which can be solved for K_2 and K_3 . The values for K_2 and K_3 obtained were constant in most cases. It must be understood that the work is still in progress and that the values for V_1 and V_2 which are used in the above are not the actual values. It is seen that the values $K_2 = 1.70$ and

$K_e = -40$ as averages for the values thus obtained, we can then substitute them in the above equation and calculate our values for K_n' which we should obtain experimentally if the equation holds. The table on page 62 shows the calculated and found values with the percentage error.

$\bar{V}_2 = -20$ as examples for the values thus obtained, we can then calculate them in the above equation and calculate the values for $\bar{V}_1$ which is shown in column 1 of Table II. The equation below. The table of page 12 shows the calculated and found values with the percentage error.

EXPERIMENTAL

Preparation of materials. The method used in preparing the solutions and carrying out the reactions at 35° is described in detail by Robertson¹.

The methyl iodide used was of special purity obtained from Hoffmann and Kruff. To insure a large yield of a constant boiling material, two kilograms of the alkyl halide were shaken with dilute sodium carbonate solution to remove any traces of hydriodic acid, washed twice with water, and then shaken for a few minutes with anhydrous sodium sulphate, and filtered through glass wool. A considerable quantity of phosphorus pentoxide was added and allowed to remain over night, after which the methyl iodide was distilled. This treatment with the phosphorus pentoxide was repeated, and the product thus obtained was fractionated. Special care was taken to have the condenser and all apparatus perfectly dry, and, with these precautions, a product was obtained, the total quantity of which boiled within a few hundredths of a degree. A table is given below showing the boiling points of the different fractions, the barometric pressure and the boiling points at 760 millimeters after all the necessary corrections are made.

1. I.C.

EXPERIMENTAL

Experiments in distillation. The method used in separating the solutions and carrying out the reactions at 40° is described in detail by Robertson.

The catalyst used was of special purity obtained from Williams and Davis. To secure a large yield of a pure product colling material, two kilograms of the catalyst were mixed with dilute sodium carbonate solution to remove any traces of phosphoric acid, washed twice with water, and then washed for a few minutes with anhydrous sodium carbonate and filtered through glass wool. A considerable quantity of phosphoric pentoxide was added and allowed to remain over night, after which the catalyst was distilled. This treatment with the phosphoric pentoxide was repeated, and the product thus obtained was fractionated. Special care was taken to have the condenser and all apparatus perfectly dry, and with these precautions, a product was obtained, the total quantity of which fell within a few hundredths of a gram. A table is given below showing the boiling points of the different fractions, the percentage pressure and the boiling points at 760 millimeters after all the necessary corrections have been made.

Observed B.pt.	Barometer	Corrected B.pt.
43.80 - 43.83°	769.4 mm.	43.17 - 43.20°
43.83 - 43.82	769.3	43.19 - 43.20
43.82 - 43.81	769.3	43.30 - 43.19
43.81 - 43.81	769.3	43.19 - 43.19
43.93 - 43.91	771.3	43.31 - 43.31
43.91 - 43.86	770.6	43.31 - 43.31
43.86 - 43.82	769.5	43.31 - 43.30
43.82 - 43.80	768.9	43.30 - 43.30
43.80 - 43.78	768.3	43.30 - 43.33
43.78 - 43.79	767.7	43.33 - 43.33

The rise in boiling point observed in the last fraction was probably due to superheating. No correction was necessary for the exposed stem because the distilling flask with a long neck made especially for the purpose was used, which made it possible to surround the total column of mercury with the vapor of the alkyl halide. In correcting for the atmospheric pressure the value used was 0.041° per millimeter. The total amount of methyl iodide recovered in these ten fractions was 1772 grams out of the 3000 grams used. This is a very satisfactory yield for such a volatile substance. In order to decrease the loss due to the volatility of the methyl iodide, ice-water was continually run through the condenser by means of a fine lead coil of several turns surrounded by ice, and the receiving bottles^{were} cooled by ice-water and fairly tightly stoppered with cotton. It was not usual for the boiling point to become constant at the very beginning of the distillation,

as did the above sample.

From another sample of methyl iodide from the same source which had been treated in a similar manner but without the special precautions taken for excluding moisture from all apparatus and for preventing the loss of methyl iodide, only about 1350 grams were obtained. This had a boiling point of $43.19 - 43.23^{\circ}$, most of which, however, came over at $43.30 - 43.33^{\circ}$ which is identically the same boiling point obtained in the sample mentioned above. The lower yield was partly due to the fact that the methyl iodide was digested twice for nearly a whole day on a water bath in the presence of phosphorus pentoxide, and it was impossible to prevent entirely the escape of the vapor from the connections and also from the top of the condenser. It was afterwards found that this heating was entirely superfluous as a sample which simply stood over night with phosphorus pentoxide gave a very satisfactory boiling point.

Another sample from Hoffmann and Kruff, treated in a similar way but without the special precautions, and distilled with the mercury column exposed, gave the corrected boiling point of $43.15 - 43.19^{\circ}$. There is no doubt in the author's mind that this discrepancy is principally the result of errors in correcting the boiling point and that the sample was practically as good as those mentioned above.

One sample of good methyl iodide which had stood during the summer was distilled once from phosphorus pentoxide and then fractionated; it gave a large middle portion which boiled at $43.31 - 43.34^{\circ}$. Another sample of methyl iodide

as all the above samples.

From various samples of heavy lithic from the same source which had been treated in a similar manner but without the special precautions taken for excluding extraneous iron, the specimens and for representing the issue of lithic lithic, only about 1000 grains were obtained. This had a boiling point of 41.15 - 41.25°, none of which, however, came over at 41.15 - 41.25° which is identical with the same boiling point obtained in the same is mentioned above. The lower yield was lithic due to the fact that the lithic lithic was distilled water for nearly a whole day on a water bath in the presence of phosphoric anhydride, and it was impossible to prevent entirely the escape of the water from the connections and also from the top of the condenser. It was afterwards found that this boiling was entirely excluded as a sample which slightly more than 40% of phosphoric anhydride gave a very satisfactory boiling point.

Another sample from Hoffman and Trill, treated in a similar way but without the special precautions, and distilled with the known column exposed, gave the corrected boiling point of 41.15 - 41.25°. There is no doubt in the author's mind that this discrepancy is principally the result of error in the method of boiling point and that the same was practically as good as those mentioned above.

The sample of heavy lithic which has been treated in the same way as mentioned above with phosphoric anhydride and then distilled, it gave a lower yield of lithic boiling at 41.15 - 41.25°. Another sample of heavy lithic

which was made in the laboratory by the method described by Robertson, gave a fraction boiling at 43.17 - 43.23°.

It would seem from the boiling points obtained that the methyl iodide used in the reactions was of especial purity. Nevertheless, it was not only a wise precaution but a point of great interest to determine the composition of such samples by analysis.

Analysis of Methyl Iodide. At first an attempt was made to use small glass bulbs which were filled with the sample, sealed, weighed, and then broken by means of a glass rod in an excess of a 2/N solution of sodium ethylate contained in a small flask, the latter being afterward corked and warmed at about 60°. After the reaction was complete the solution was acidified and the iodine was precipitated as silver iodide.

AgI obtained	AgI calculated	Per cent. error.
1.0336 gms.	1.0364 gms.	-0.3
2.1841 "	2.1938 "	-0.5

Either loss of alkyl halide occurred during the heating, or moisture was absorbed by the methyl iodide while the bulb was being filled.

Much more satisfactory results were obtained by using small weighing tubes with ground glass stoppers which were especially made for the purpose and held about two cubic centimeters. The alkyl halide could thus be weighed accurately and without loss. The weighing tube containing the alkyl halide was then placed in a regular Carius tube containing an excess of sodium ethylate, sealed, and heated a sufficient

length of time in a water bath. It was necessary, of course, to mix the contents thoroughly before heating. The stopper in the little weighing tube was easily removed by warming the tube slightly and then shaking it gently. When the reaction was complete the contents of the tube were washed into a beaker, and then filtered into a calibrated flask, diluted to the mark and weighed. Each portion taken out for analysis was then weighed, since this procedure was much more accurate than the volumetric method.

Sample I.

AgI obtained	AgI calculated	Per cent. error.
2.0346 gms.	2.0327 gms.	+0.1
2.0336	2.0339	-0.014
1.9956	1.9953	+0.015

Sample II.

A

AgI obtained	AgI calculated	Per cent. error.
1.5890 gms.	1.5889 gms.	0.00
1.5886	1.5889	-0.03
1.5894	1.5898	-0.035

B

1.7333 gms.	1.7343 gms.	-0.05
1.7364	1.7380	-0.10
1.7371	1.7373	-0.013

C

1.5493 gms.	1.5500 gms.	-0.05
1.5469	1.5495	-0.04
1.5486	1.5499	-0.07

In the above analyses Sample I and Sample II were taken from different samples of methyl iodide.

It is seen that we can easily make alkyl halides with a high degree of purity and that this method of analysis is very accurate.

Preparation of solutions for salt catalysis. The alcohol used contained from 0.03 - 0.04 per cent. of water. The preparation of this alcohol has been previously described by Robertson as was also the regulation of the bath at 25°, and the mode of procedure used in determining the velocity constants at this temperature.

The solutions for the experiments on the salt effect were usually prepared by making a sodium ethylate solution having a strength slightly greater than that desired. The calculated number of cubic centimeters of the solution, say 90 - 95 c.c., were put into a 100 c.c. flask which already contained the amount of sodium iodide required, diluted to the mark, and then 10 c.c. of this was titrated to obtain the value of A. A very small excess of the ethylate solution was usually added to be sure that the resulting solution was not too dilute. If too concentrated, the calculated amount of alcohol was added; if this was more than a few tenths of one cubic centimeter, more sodium iodide was added as well. The amount of salt added in these cases was too small to introduce any appreciable error in the volume.

Calibration of the machine. Since methyl iodide is so very volatile it would be impossible to transfer the solution to the reaction flasks by the use of an ordinary pipette, without losing a large amount of the alkyl halide. A special ap-

paratus called the "machine" was devised to transfer these solutions with the least possible loss. This machine was described in detail by Robertson.

The machine pipette was first marked to deliver 10 c.c. of absolute alcohol at 20° and the volume obtained was 10.004 c.c., the error being less than 0.1 per cent. Since this pipette was to be used for alkyl halide solutions, the author wished to determine whether the same volume of methyl iodide solution would be obtained. The specific gravity of the methyl iodide solution was determined and successive portions of 10 c.c. each were then run out into the same small flasks used for reactions, and weighed. These flasks were closed by corks but it had been shown previously that the reaction mixture did not escape, and in this case it was found that these flasks containing the volatile methyl iodide solution could be weighed without loss. The volumes of N/2 and N/4 alkyl halide solutions obtained at 25° were 10.00, 9.997, 9.982, 9.992, 9.996, and 9.990 c.c. The average of these is 9.993 c.c. and was considered as sufficiently near to the desired volume. Although this difference is probably within the experimental error, yet it is quite possible that the alkyl halide solution is slightly more viscous and does not drain as readily.

When using this same machine at 0° the pipette was recalibrated with a methyl iodide solution at this temperature. The volumes obtained were 9.979, 9.981, 9.971, 9.975, and 9.974 c.c. the average being 9.976 c.c. In all of the work at 0° the solution was run out to a point on the pipette which would deliver exactly 10 c.c. at this temperature.

Determination of loss of alkyl halide. An investigation was also carried out to determine whether it was necessary of the machine to keep the tip of the delivery tube/below the surface of the liquid in the reaction flask while introducing the alkyl halide solution.

AgI obtained	AgI calculated.	Strength of solution	Percent-age error.	Position of tip
2.3483	2.3550	(100.3 o/o)	-0.38	Below surface of liquid.
2.3374	2.3550	"	-0.75	Just below neck of flask.
2.3330	2.3550	"	-0.90	Near top of neck.
2.3455	2.3550	"	-0.40	Just below neck.
2.3330	2.3550	"	-0.90	Half way down neck.

These figures show that it is certainly necessary to keep the tip well under the surface of the liquid.

A large number of determinations were also carried out to see whether the three tenths per cent. excess of methyl iodide ordinarily added was sufficient to make up for any loss of the alkyl halide during the preparation and transfer of the solution. While keeping the conditions the same as those used for studying the reactions, 10 c.c. of the solution was run into a small reaction flask containing a 50 per cent. alcoholic solution of silver nitrate. The results obtained are given in the tables below. The calculated values given for the silver iodide include the excess added which is given in the parenthesis under "Strength of solution".

AgI obtained	AgI calculated	Strength of solution	Per cent. error.
3.3535 gms.	3.3550 gms.	(100.10/o)	-0.06
3.3500	3.3550	(100.1 ")	-0.21
2.3424	2.3550	(100.1 ")	-0.11
2.3570	2.3620	(100.6 ")	-0.21
2.3533	2.3597	(100.5 ")	-0.27
1.1727	1.1775	(100.3 ")	-0.40
1.1713	1.1775	(100.3 ")	-0.53
			Average = -0.36

At 0°.

AgI obtained	AgI calculated	Strength of solution	Per cent. error.
1.1743*gms.	1.1799 gms.	(100.50/o)	-0.47
1.1729*	1.1799	(100.5 ")	-0.60
1.1730*	1.1799	(100.5 ")	-0.60
2.3584	2.3598	(100.5 ")	-0.06
4.7279	4.7288	(100.7 ")	-0.02
2.3576	2.3621	(100.6 ")	-0.20
2.3583	2.3621	(100.6 ")	-0.16
2.3542	2.3598	(100.5 ")	-0.22
4.7057	4.7101	(100.3 ")	-0.09
2.3495	2.3598	(100.5 ")	-0.43
* Same solution			
			Average = -0.37

The average of the above analyses shows that the 0.3 per cent. excess of methyl iodide seems to be sufficient although there is considerable variation in the results. However, in a number of reactions 0.5 per cent. excess was added.

Year	Age	Age	Age
1900	100.0	100.0	100.0
1901	100.0	100.0	100.0
1902	100.0	100.0	100.0
1903	100.0	100.0	100.0
1904	100.0	100.0	100.0
1905	100.0	100.0	100.0
1906	100.0	100.0	100.0
1907	100.0	100.0	100.0
1908	100.0	100.0	100.0
1909	100.0	100.0	100.0
1910	100.0	100.0	100.0
1911	100.0	100.0	100.0
1912	100.0	100.0	100.0
1913	100.0	100.0	100.0
1914	100.0	100.0	100.0
1915	100.0	100.0	100.0
1916	100.0	100.0	100.0
1917	100.0	100.0	100.0
1918	100.0	100.0	100.0
1919	100.0	100.0	100.0
1920	100.0	100.0	100.0

1920

Year	Age	Age	Age
1921	100.0	100.0	100.0
1922	100.0	100.0	100.0
1923	100.0	100.0	100.0
1924	100.0	100.0	100.0
1925	100.0	100.0	100.0
1926	100.0	100.0	100.0
1927	100.0	100.0	100.0
1928	100.0	100.0	100.0
1929	100.0	100.0	100.0
1930	100.0	100.0	100.0
1931	100.0	100.0	100.0
1932	100.0	100.0	100.0
1933	100.0	100.0	100.0
1934	100.0	100.0	100.0
1935	100.0	100.0	100.0
1936	100.0	100.0	100.0
1937	100.0	100.0	100.0
1938	100.0	100.0	100.0
1939	100.0	100.0	100.0
1940	100.0	100.0	100.0
1941	100.0	100.0	100.0
1942	100.0	100.0	100.0
1943	100.0	100.0	100.0
1944	100.0	100.0	100.0
1945	100.0	100.0	100.0
1946	100.0	100.0	100.0
1947	100.0	100.0	100.0
1948	100.0	100.0	100.0
1949	100.0	100.0	100.0
1950	100.0	100.0	100.0

The number of the above employees who had been in the service of the company for a period of ten years or more, as of the end of the year 1950, is shown in the following table. It is noted that the number of employees who had been in the service of the company for a period of ten years or more, as of the end of the year 1950, is 1,000.

Reaction velocities at 0°. For the study of the velocity of the reaction between methyl iodide, sodium, lithium, and potassium ethylates at 0° an ice bath was used which gave very satisfactory results. This consisted of a copper bath ten inches deep and fourteen inches in diameter which was tinned on the inside, and placed in a wooden box containing lamb's wool. The bottom was protected with a thick felt pad about one inch thick, and the top, by a tightly fitting metal cover over which were placed several pads of felt and wool. The metal cover was made in two pieces, one of which was removed when it was necessary to take out the flask, start the reaction, etc. The bath as set up ready for use is shown in the diagram on page 64. Special attention is called to the stirring device. The motor is supported by an iron framework which is bolted firmly to the wall to prevent vibration. Motion is imparted to the propellor shaft by a friction pulley, by means of which the speed can be easily controlled.

Tests with a Beckmann thermometer showed that all parts of the bath remained within a few thousandths of a degree of zero. The ethylate solutions were made up at 0°, and transferred at this temperature into the reaction flasks by surrounding the pipette with a jacket containing cracked ice and water. In beginning this work the jacket of the machine pipette was filled only once or twice with ice water during the seven or eight minutes required for running out all of the alkyl halide solution. The reaction flask containing the ethylate solution had to be lifted from the ice bath in order to keep

the delivery tip of the machine under the surface of the liquid in the reaction flask. The tube was then shaken by inverting it a few times, and then put back into the bath. Although this process required only one minute for each tube, test with a thermometer showed that the temperature had risen several degrees, and that some better method had to be devised. The possibility of mixing the solution under ice water was suggested, but as this method would have required new apparatus, the following method was tried with very satisfactory results. Ice water from the bath was run continuously through the jacket of the machine by means of a pump run by a small electric motor. Although the exposed circulation was made as short as possible, the temperature of the water returning to the bath had risen several degrees. Even when this current of water was returned to the bath in the vicinity of the thermometer, no change in the temperature could be detected unless the water played directly upon the thermometer. In that part of the bath which contained the reaction flasks, no change in temperature could be detected. The delivery tube of the machine and all exposed connecting tubes were protected by several layers of binding tape, and that liquid in the most exposed part of the delivery tip was discarded just before each 10 c.c. portion was run out. The reaction flask containing the ethylate solution was lifted from the bath in a beaker containing cracked ice and water, and after the alkyl halide solution had been transferred to it, the flask was returned to the bath as rapidly as possible. Tests made with the thermometer showed that by this method the temperature was kept below 0.5° , and the zero point was quickly

The analysis of the results of the investigation of the
 liquid in the reaction vessel, the water and the water
 of investigation is a few times, and then the water is
 only. Although this process is not very accurate, but
 each time, just with a thermometer showed that the tempera-
 ture had risen several degrees, and then some water was
 added to be removed. The possibility of this is not
 taken into account, and as this is not a very accurate
 method, the following method was used: the
 very satisfactory results. The water from the water was
 continuously through the liquid of the reaction by means of
 a few mm of a water electric motor. Although the energy
 situation was made as short as possible, the temperature
 of the water resulting to the water was several degrees.
 From when this amount of water was added to the water in
 the vicinity of the reaction, it showed in the instrument
 could be detected unless the water liquid is not very
 important. In that case of the water which contained the re-
 action liquid, as shown in temperature could be detected. The
 addition of the water and the water was not necessary, because
 the addition of several degrees of water, and the
 liquid in the water, and of the liquid in the water
 added. And before each 10.0. addition was made. The re-
 action liquid, according to the water reaction was added from
 the water in a certain temperature, and the water was added
 for the liquid and the water was added to it. The
 liquid was removed to the water in a certain temperature. The
 water with the liquid in the water was added to the water
 liquid, and the water was added to the water.

reached by shaking the flask two minutes in the bath. For the work with the N/16 and N/32 solutions 40 c.c. portions were used instead of the usual 30 c.c., and the solutions were mixed by inverting and shaking the flasks under the ice water. The corks were protected by "rubber fingers" over the necks of the flasks. A rise in the temperature of even several tenths of a degree for one minute would have but little effect on the constants, especially those for the dilute solutions for which the time periods were so long.

Lithium Ethylate and Methyl Iodide at 0°C.

Table I

0.5 N Lithium Ethylate and
0.5 Methyl Iodide at 0°C.

$$A = 9.98$$

t	x	K_v
180	1.782	0.001207
340	2.226	0.001196
300	2.642	0.001200
382	3.156	0.001210
1378	5.946	0.001153
1667	6.580	0.001161

$$\text{Average} = 0.001188$$

$$K_n = 0.002381$$

$$K_n^* = 0.002411$$

Table II

0.5 N Lithium Ethylate and
0.5 N Methyl Iodide at 0°C.

$$A = 9.98$$

t	x	K_v
205	2.016	0.001234
266	2.448	0.001233
347	2.972	0.001232
1353	6.120	0.001172
1430	6.264	0.001180

$$\text{Average} = 0.001206$$

$$K_n = 0.002417$$

$$K_n^* = 0.002425$$

Table III

0.5 N Lithium Ethylate and
1.0 N Methyl Iodide at 0°C.

$$A = 10.00 \quad B = 30.00$$

t	x	K_v
180	3.318	0.001232
356	4.240	0.001234
330	4.960	0.001214
424	5.690	0.001195
520	6.304	0.001186

$$\text{Average} = 0.001210$$

$$K_n = 0.002420$$

Table IV

0.25 N Lithium Ethylate and
0.5 N Methyl Iodide at 0°C.

$$A = 10.01 \quad B = 30.00$$

t	x	K_v
142	1.722	0.0006967
197	2.246	0.0006863
259	2.790	0.0006826
318	3.250	0.0006748
348	3.470	0.0006739

$$\text{Average} = 0.0006828$$

$$K_n = 0.002728$$

* If last two values of K_v are omitted.

1.1.1.1

THE UNIVERSITY OF CHICAGO LIBRARY

THE UNIVERSITY OF CHICAGO LIBRARY

THE UNIVERSITY OF CHICAGO LIBRARY

THE UNIVERSITY OF CHICAGO LIBRARY

THE UNIVERSITY OF CHICAGO LIBRARY

THE UNIVERSITY OF CHICAGO LIBRARY

THE UNIVERSITY OF CHICAGO LIBRARY

1.1.1.2

THE UNIVERSITY OF CHICAGO LIBRARY

THE UNIVERSITY OF CHICAGO LIBRARY

THE UNIVERSITY OF CHICAGO LIBRARY

THE UNIVERSITY OF CHICAGO LIBRARY

THE UNIVERSITY OF CHICAGO LIBRARY

THE UNIVERSITY OF CHICAGO LIBRARY

1.1.1.3

THE UNIVERSITY OF CHICAGO LIBRARY

THE UNIVERSITY OF CHICAGO LIBRARY

THE UNIVERSITY OF CHICAGO LIBRARY

THE UNIVERSITY OF CHICAGO LIBRARY

THE UNIVERSITY OF CHICAGO LIBRARY

THE UNIVERSITY OF CHICAGO LIBRARY

THE UNIVERSITY OF CHICAGO LIBRARY

1.1.1.4

THE UNIVERSITY OF CHICAGO LIBRARY

THE UNIVERSITY OF CHICAGO LIBRARY

THE UNIVERSITY OF CHICAGO LIBRARY

THE UNIVERSITY OF CHICAGO LIBRARY

THE UNIVERSITY OF CHICAGO LIBRARY

THE UNIVERSITY OF CHICAGO LIBRARY

Table V

0.25 N Lithium Ethylate and
0.5 N Methyl Iodide at 0°C.

A = 10.03 B = 30.00

t	x	K_v
344	3.456	0.0006805
433	3.976	0.0006730
502	4.460	0.0006730
1480	7.613	0.0006420
1608	7.830	0.0006367

Average = 0.0006567

$K_n = 0.003633$

$K_n^* = 0.003696$

Table VI

0.35 N Lithium Ethylate and
1.0 N Methyl Iodide at 0°C.

A = 10.03 B = 40.00

t	x	K_v
187	3.840	0.0006831
230	4.444	0.0006803
300	5.256	0.0006714
340	5.656	0.0006670
390	6.104	0.0006630
440	6.500	0.0006597

Average = 0.0006707

$K_n = 0.003678$

Table VII

0.135 N Lithium Ethylate and
0.35 N Methyl Iodide at 0°C.

A = 10.04 B = 30.00

t	x	K_v
1016	4.832	0.0003767
1077	4.992	0.0003746
1140	5.144	0.0003719
1201	5.304	0.0003718
1277	5.488	0.0003710
1347	5.698	0.0003712

Average = 0.0003729

$K_n = 0.003971$

Table VIII

0.135 N Lithium Ethylate and
0.1373 N Methyl Iodide at 0°C.

A = 10.00 B = 10.18

t	x	K_v
1098	2.960	0.0003745
1194	3.136	0.0003742
1289	3.296	0.0003730
1428	3.636	0.0003743
1509	3.656	0.0003732
2619	4.992	0.0003705

Average = 0.0003733

$K_n = 0.003986$

* If last two values of K_v are omitted.

TABLE 1

For $\alpha = 0.05$ and $\beta = 0.1$
 $\alpha = 0.05$ $\beta = 0.1$

$\sqrt{n}$	α	β
100000.0	0.05	0.1
100000.0	0.05	0.1
100000.0	0.05	0.1
100000.0	0.05	0.1
100000.0	0.05	0.1
100000.0	0.05	0.1

100000.0 = 0.05

100000.0 = 0.1

TABLE 2

For $\alpha = 0.05$ and $\beta = 0.1$
 $\alpha = 0.05$ $\beta = 0.1$

$\sqrt{n}$	α	β
100000.0	0.05	0.1
100000.0	0.05	0.1
100000.0	0.05	0.1
100000.0	0.05	0.1
100000.0	0.05	0.1
100000.0	0.05	0.1

100000.0 = 0.05

100000.0 = 0.1

100000.0 = 0.1

TABLE 3

For $\alpha = 0.05$ and $\beta = 0.1$
 $\alpha = 0.05$ $\beta = 0.1$

$\sqrt{n}$	α	β
100000.0	0.05	0.1
100000.0	0.05	0.1
100000.0	0.05	0.1
100000.0	0.05	0.1
100000.0	0.05	0.1
100000.0	0.05	0.1

100000.0 = 0.05

100000.0 = 0.1

TABLE 4

For $\alpha = 0.05$ and $\beta = 0.1$
 $\alpha = 0.05$ $\beta = 0.1$

$\sqrt{n}$	α	β
100000.0	0.05	0.1
100000.0	0.05	0.1
100000.0	0.05	0.1
100000.0	0.05	0.1
100000.0	0.05	0.1
100000.0	0.05	0.1

100000.0 = 0.05

100000.0 = 0.1

Table IX

0.0625 N Lithium Ethylate and
0.35 N Methyl Iodide at 0°C.

$$A = 9.99 \quad B = 40.00$$

t	x	K_v
585	3.706	0.0003084
657	4.030	0.0003078
830	4.758	0.0003085
905	4.994	0.0003058
1034	5.433	0.0003050
3108	7.750	0.0003021

$$\text{Average} = 0.0003063$$

$$K_n = 0.003304$$

Table X

0.0625 N Lithium Ethylate and
0.1289 N Methyl Iodide at 0°C.

$$A = 10.01 \quad B = 30.63$$

t	x	K_v
976	3.198	0.0003144
1066	3.394	0.0003125
1134	3.570	0.0003140
1370	3.858	0.0003130
1378	4.086	0.0003133
2514	6.058	0.0003234
3777	6.154	0.0003088

$$\text{Average} = 0.0003142$$

$$K_n = 0.003433$$

Table XI

0.0625 N Lithium Ethylate and
0.35 N Methyl Iodide at 0°C.

$$A=10.00 \quad B=40.00$$

t	x	K_v
567	3.640	0.0003098
612	3.840	0.0003088
880	4.140	0.0003083
780	4.540	0.0003070
900	5.008	0.0003077
1020	5.416	0.0003072

$$\text{Average} = 0.0003081$$

$$K_n = 0.003330$$

Table XII

0.0625 N Lithium Ethylate and
0.35 N Methyl Iodide at 0°C.

$$A = 10.02 \quad B = 40.00$$

t	x	K_v
1045	5.444	0.0003036
1135	5.692	0.0003019
1202	5.934	0.0003041
1326	6.244	0.0003031
1496	6.628	0.0003014
3480	8.212	0.0001997

$$\text{Average} = 0.0003023$$

$$K_n = 0.003231$$

TABLE 1

TABLE 1. Values of the function $f(x)$ for $x = 0.1, 0.2, \dots, 1.0$. The function is defined by the formula $f(x) = x^2 + 1$.

x	$f(x)$
0.1	1.01
0.2	1.04
0.3	1.09
0.4	1.16
0.5	1.25
0.6	1.36
0.7	1.49
0.8	1.64
0.9	1.81
1.0	2.00

TABLE 1. Values of the function $f(x)$ for $x = 0.1, 0.2, \dots, 1.0$.

TABLE 1. Values of the function $f(x)$ for $x = 0.1, 0.2, \dots, 1.0$.

TABLE 2

TABLE 2. Values of the function $g(x)$ for $x = 0.1, 0.2, \dots, 1.0$. The function is defined by the formula $g(x) = x^2 - 1$.

x	$g(x)$
0.1	-0.99
0.2	-0.96
0.3	-0.91
0.4	-0.84
0.5	-0.75
0.6	-0.64
0.7	-0.51
0.8	-0.36
0.9	-0.19
1.0	0.00

TABLE 2. Values of the function $g(x)$ for $x = 0.1, 0.2, \dots, 1.0$.

TABLE 2. Values of the function $g(x)$ for $x = 0.1, 0.2, \dots, 1.0$.

TABLE 3

TABLE 3. Values of the function $h(x)$ for $x = 0.1, 0.2, \dots, 1.0$. The function is defined by the formula $h(x) = x^2 + 2$.

x	$h(x)$
0.1	2.01
0.2	2.04
0.3	2.09
0.4	2.16
0.5	2.25
0.6	2.36
0.7	2.49
0.8	2.64
0.9	2.81
1.0	3.00

TABLE 3. Values of the function $h(x)$ for $x = 0.1, 0.2, \dots, 1.0$.

TABLE 3. Values of the function $h(x)$ for $x = 0.1, 0.2, \dots, 1.0$.

TABLE 4

TABLE 4. Values of the function $k(x)$ for $x = 0.1, 0.2, \dots, 1.0$. The function is defined by the formula $k(x) = x^2 - 2$.

x	$k(x)$
0.1	-1.99
0.2	-1.96
0.3	-1.91
0.4	-1.84
0.5	-1.75
0.6	-1.64
0.7	-1.51
0.8	-1.36
0.9	-1.19
1.0	-1.00

TABLE 4. Values of the function $k(x)$ for $x = 0.1, 0.2, \dots, 1.0$.

TABLE 4. Values of the function $k(x)$ for $x = 0.1, 0.2, \dots, 1.0$.

Table XIII

0.03135 N Lithium Ethylate and
0.35 N Methyl Iodide at 0°C.

A = 10.03 B = 80.00

t	x	K_v
570	4.030	0.0001162
630	4.326	0.0001157
690	4.590	0.0001148
760	4.934	0.0001156
890	5.430	0.0001142
1080	6.070	0.0001129

Average = 0.0001149

K_n = 0.003665

Table XIV

0.03125 N Lithium Ethylate and
0.135 N Methyl Iodide at 0°C.

A = 10.00 B = 40.00

t	x	K_v
1067	3.752	0.0001161
1136	3.680	0.0001141
1287	4.312	0.0001166
1463	4.634	0.0001134
2423	6.388	0.0001138
3896	6.880	0.0001136

Average = 0.0001143

K_n = 0.003657

Table XV

0.03135 N Lithium Ethylate and
0.35 N Methyl Iodide at 0°C.

A = 10.00 B = 80.00

t	x	K_v
561	3.840	0.0001108
647	4.328	0.0001129
816	5.000	0.0001100
898	5.296	0.0001090
983	5.664	0.0001107
1053	5.938	0.0001114
2086	8.168	0.0001087

Average = 0.0001105

K_n = 0.003536

TABLE 1

See Table 1, Column 1, for details.
 The following table shows the
 results of the analysis.

λ	μ	ν
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000

TABLE 1, Column 1, for details.

TABLE 1, Column 1, for details.

TABLE 2

See Table 2, Column 1, for details.
 The following table shows the
 results of the analysis.

λ	μ	ν
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000

TABLE 2, Column 1, for details.

TABLE 2, Column 1, for details.

TABLE 3

See Table 3, Column 1, for details.
 The following table shows the
 results of the analysis.

λ	μ	ν
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000

TABLE 3, Column 1, for details.

TABLE 3, Column 1, for details.

Table 1. Summary of the results of the analysis of variance for the different factors.

Factor	df	SS	MS	F	P
Block	1	10.0	10.0	1.0	0.3
Replication	1	10.0	10.0	1.0	0.3
Treatment	1	10.0	10.0	1.0	0.3
Error	1	10.0	10.0	1.0	0.3
Total	4	40.0	10.0		
Block	1	10.0	10.0	1.0	0.3
Replication	1	10.0	10.0	1.0	0.3
Treatment	1	10.0	10.0	1.0	0.3
Error	1	10.0	10.0	1.0	0.3
Total	4	40.0	10.0		
Block	1	10.0	10.0	1.0	0.3
Replication	1	10.0	10.0	1.0	0.3
Treatment	1	10.0	10.0	1.0	0.3
Error	1	10.0	10.0	1.0	0.3
Total	4	40.0	10.0		

Table 2. Summary of the results of the analysis of variance for the different factors.

Table 2

Table 3. Summary of the results of the analysis of variance for the different factors.

Factor	df	SS	MS	F	P
Block	1	10.0	10.0	1.0	0.3
Replication	1	10.0	10.0	1.0	0.3
Treatment	1	10.0	10.0	1.0	0.3
Error	1	10.0	10.0	1.0	0.3
Total	4	40.0	10.0		
Block	1	10.0	10.0	1.0	0.3
Replication	1	10.0	10.0	1.0	0.3
Treatment	1	10.0	10.0	1.0	0.3
Error	1	10.0	10.0	1.0	0.3
Total	4	40.0	10.0		

Table 4. Summary of the results of the analysis of variance for the different factors.

Table XVI

Summary of Lithium Ethylate and Methyl Iodide at 0°C.

V	K_n	Average K_n
2	0.003430 0.003381 (0.003411) 0.003417 (0.003435)	0.003406 (0.003419)
4	0.002728 0.002678 0.002663 (0.002696)	0.002690 (0.002700)
8	0.002971 0.002986	0.002978
16	0.003304 0.003423 0.003330 0.003331	0.003323
32	0.003665 0.003657 0.003536	0.003619

Values in parenthesis under K_n refer to starred values under tables average K_n in parenthesis, only to the starred values in these tables

Table XVII

Lithium Ethylate and Methyl Iodide at 0°C.

V	α	$1 - \alpha$	K_n found		K_n calculated		Per cent. e
			a	b	a	b	a
2	0.144	0.856	0.002406	0.002419	0.002440	0.002450	-1.40 -1
4	0.230	0.730	0.002690	0.002700	0.002690	0.002697	0.0 +0
8	0.312	0.688	0.002978	0.002978	0.002991	0.002994	-0.43 -0
16	0.409	0.591	0.003322	0.003323	0.003310	0.003308	+0.36 +0
32	0.519	0.481	0.003619	0.003619	0.003668	0.003662	-1.30 -1

Values under "b" are calculated on values of K_n in parenthesis.

Table XVIII

K_1 and K_m for Lithium Ethylate and Methyl Iodide at 0°C.

V	K_1		K_m	
	a	b	a	b
2 : 4	0.005590	0.005569	0.001872	0.001891
3 : 8	0.005317	0.005263	0.001918	0.001943
2 : 16	0.005360	0.005331	0.001911	0.001931
2 : 32	0.005174	0.005157	0.001943	0.001960
4 : 8	0.005134	0.005060	0.002001	0.002035
4 : 16	0.005297	0.005264	0.001955	0.001977
4 : 32	0.005115	0.005098	0.002006	0.002024
8 : 16	0.005411	0.005411	0.001875	0.001875
8 : 32	0.005109	0.005109	0.002012	0.002012
16 : 32	0.004922	0.004922	0.002214	0.002314
Average = 0.005243 0.005218 0.001971 0.001986				

Values under "b" are calculated from values of Average K_n in parenthesis.

TITLE PAGE

THE 1980-1981 FISH AND WILDLIFE SURVEY OF THE GREAT LAKES

1	2	3
1. NAME OF SURVEY	2. NAME OF SURVEY	3. DATE
4. NAME OF SURVEY	5. NAME OF SURVEY	6. DATE
7. NAME OF SURVEY	8. NAME OF SURVEY	9. DATE
10. NAME OF SURVEY	11. NAME OF SURVEY	12. DATE
13. NAME OF SURVEY	14. NAME OF SURVEY	15. DATE
16. NAME OF SURVEY	17. NAME OF SURVEY	18. DATE
19. NAME OF SURVEY	20. NAME OF SURVEY	21. DATE
22. NAME OF SURVEY	23. NAME OF SURVEY	24. DATE
25. NAME OF SURVEY	26. NAME OF SURVEY	27. DATE
28. NAME OF SURVEY	29. NAME OF SURVEY	30. DATE
31. NAME OF SURVEY	32. NAME OF SURVEY	33. DATE
34. NAME OF SURVEY	35. NAME OF SURVEY	36. DATE
37. NAME OF SURVEY	38. NAME OF SURVEY	39. DATE
40. NAME OF SURVEY	41. NAME OF SURVEY	42. DATE
43. NAME OF SURVEY	44. NAME OF SURVEY	45. DATE
46. NAME OF SURVEY	47. NAME OF SURVEY	48. DATE
49. NAME OF SURVEY	50. NAME OF SURVEY	51. DATE
52. NAME OF SURVEY	53. NAME OF SURVEY	54. DATE
55. NAME OF SURVEY	56. NAME OF SURVEY	57. DATE
58. NAME OF SURVEY	59. NAME OF SURVEY	60. DATE
61. NAME OF SURVEY	62. NAME OF SURVEY	63. DATE
64. NAME OF SURVEY	65. NAME OF SURVEY	66. DATE
67. NAME OF SURVEY	68. NAME OF SURVEY	69. DATE
70. NAME OF SURVEY	71. NAME OF SURVEY	72. DATE
73. NAME OF SURVEY	74. NAME OF SURVEY	75. DATE
76. NAME OF SURVEY	77. NAME OF SURVEY	78. DATE
79. NAME OF SURVEY	80. NAME OF SURVEY	79. DATE
82. NAME OF SURVEY	83. NAME OF SURVEY	82. DATE
85. NAME OF SURVEY	86. NAME OF SURVEY	85. DATE
88. NAME OF SURVEY	89. NAME OF SURVEY	88. DATE
91. NAME OF SURVEY	92. NAME OF SURVEY	91. DATE
94. NAME OF SURVEY	95. NAME OF SURVEY	94. DATE
97. NAME OF SURVEY	98. NAME OF SURVEY	97. DATE
100. NAME OF SURVEY	99. NAME OF SURVEY	100. DATE

NAME OF SURVEY: NAME OF SURVEY: NAME OF SURVEY

NAME OF SURVEY: NAME OF SURVEY: NAME OF SURVEY

Sodium Ethylate and Methyl Iodide at 0°C.

Table XIX

1.0 N Sodium Ethylate and
1.0 N Methyl Iodide at 0°C.

$$A = 9.97$$

t	x	K_v
150	3.330	0.003343
180	3.733	0.003325
210	4.088	0.003310
240	4.400	0.003291
1260	8.000	0.003223
1421	8.192	0.003244

$$\text{Average} = 0.003289$$

$$K_n = 0.003399$$

Table XX

1.0 N Sodium Ethylate and
1.0 N Methyl Iodide at 0°C.

$$A = 10.03$$

t	x	K_v
120	2.857	0.003319
150	3.319	0.003297
184	3.778	0.003284
197	3.922	0.003259
230	4.218	0.003299
1168	7.904	0.003183

$$\text{Average} = 0.003274$$

$$K_n = 0.003264$$

Table XXI

0.5 N Sodium Ethylate and
0.5 N Methyl Iodide at 0°C.

$$A = 10.00$$

t	x	K_v
150	2.058	0.001728
180	2.342	0.001700
210	2.634	0.001703
240	2.896	0.001698
2728	8.200	0.001670

$$\text{Average} = 0.001700$$

$$K_n = 0.003400$$

Table XXII

0.5 N Sodium Ethylate and
0.5 N Methyl Iodide at 0°C.

$$A = 10.03$$

t	x	K_v
180	2.372	0.001723
200	2.556	0.001712
230	2.843	0.001731
264	3.122	0.001714
290	3.342	0.001725
1230	6.770	0.001693

$$\text{Average} = 0.001715$$

$$K_n = 0.003430$$

TABLE 1. Results of the first two experiments

Experiment 1

Results of the first two experiments

Results of the first two experiments

Results of the first two experiments

λ	μ	ν
0.00000	0.00000	0.00000
0.00000	0.00000	0.00000
0.00000	0.00000	0.00000
0.00000	0.00000	0.00000
0.00000	0.00000	0.00000
0.00000	0.00000	0.00000
0.00000	0.00000	0.00000

Results of the first two experiments

Results of the first two experiments

Experiment 2

Results of the first two experiments

Results of the first two experiments

Results of the first two experiments

λ	μ	ν
0.00000	0.00000	0.00000
0.00000	0.00000	0.00000
0.00000	0.00000	0.00000
0.00000	0.00000	0.00000
0.00000	0.00000	0.00000
0.00000	0.00000	0.00000
0.00000	0.00000	0.00000

Results of the first two experiments

Results of the first two experiments

Experiment 3

Results of the first two experiments

Results of the first two experiments

Results of the first two experiments

λ	μ	ν
0.00000	0.00000	0.00000
0.00000	0.00000	0.00000
0.00000	0.00000	0.00000
0.00000	0.00000	0.00000
0.00000	0.00000	0.00000
0.00000	0.00000	0.00000
0.00000	0.00000	0.00000

Results of the first two experiments

Results of the first two experiments

Experiment 4

Results of the first two experiments

Results of the first two experiments

Results of the first two experiments

λ	μ	ν
0.00000	0.00000	0.00000
0.00000	0.00000	0.00000
0.00000	0.00000	0.00000
0.00000	0.00000	0.00000
0.00000	0.00000	0.00000
0.00000	0.00000	0.00000
0.00000	0.00000	0.00000

Results of the first two experiments

Results of the first two experiments

Table XXIII

0.5 N Sodium Ethylate and
1.0 N Methyl Iodide at 0°C.

A = 9.98 B = 30.00

t	x	K _v
100	3.782	0.001762
210	4.686	0.001740
260	5.350	0.001750
300	5.764	0.001732
364	6.366	0.001731
430	6.862	0.001721

Average = 0.001739

K_n = 0.003485

Table XXIV

0.25 N Sodium Ethylate and
0.50 N Methyl Iodide at 0°C.

A = 10.00 B = 30.00

t	x	K _v
180	2.680	0.0009337
200	2.880	0.0009207
240	3.320	0.0009245
270	3.592	0.0009148
300	3.864	0.0009121
330	4.152	0.0009203
1270	8.120	0.0009056

Average = 0.0009188

K_n = 0.003679

Table XXV

0.25 N Sodium Ethylate and
0.25 N Methyl Iodide at 0°C.

A = 10.00

t	x	K _v
180	1.400	0.0009044
210	1.608	0.0009126
250	1.840	0.0009020
280	2.020	0.0009042
301	2.132	0.0009003
333	2.328	0.0009112
1280	5.360	0.0009025

Average = 0.0009053

K_n = 0.003621

Table XXVI

0.125 N Sodium Ethylate and
0.25 N Methyl Iodide at 0°C.

A = 10.00 B = 30.00

t	x	K _v
260	2.088	0.0004765
280	2.216	0.0004751
300	2.168	0.0004806
1232	6.160	0.0004779
1326	6.384	0.0004771
1409	6.600	0.0004812

Average = 0.0004781

K_n = 0.003625

Table XXVII

0.125 N Sodium Ethylate and
0.25 N Methyl Iodide at 0°C.

A = 10.00 B = 20.00

t	x	K _v
340	1.968	0.0004813
300	2.384	0.0004816
1275	6.288	0.0004811
1360	6.480	0.0004797
1420	6.608	0.0004785
1640	7.088	0.0004852

Average = 0.0004817

K_n = 0.003854

Table XXVIII

0.0625 N Sodium Ethylate and
0.25 N Methyl Iodide at 0°C.

A = 9.96 B = 40.00

t	x	K _v
988	5.824	0.0002420
1110	6.192	0.0002400
1180	6.480	0.0002457
1265	6.752	0.0002483
1340	6.856	0.0002419
1424	7.112	0.0002458

Average = 0.0002440

K_n = 0.003919

Table XXIX

0.0625 N Sodium Ethylate and
0.25 N Methyl Iodide at 0°C.

A = 10.02 B = 40.00

t	x	K _v
360	2.900	0.0002475
390	3.140	0.0002520
420	3.340	0.0002532
1530	7.396	0.0002479
1620	7.572	0.0002473

Average = 0.0002496

K_n = 0.003985

Table XXX

0.0625 N Sodium Ethylate and
0.25 N Methyl Iodide at 0°C.

A = 10.01 B = 40.00

t	x	K _v
1106	6.418	0.0002565
1180	6.622	0.0002552
1270	6.830	0.0002521
1350	7.070	0.0002548
1447	7.278	0.0002531

Average = 0.0002543

K_n = 0.004063

TABLE 1

See Appendix Table 1 for details.

See Appendix Table 1 for details.

$$C_1 = 1, C_2 = 1$$

β	α	γ
0.00000, 0	0.00, 0	0.00
0.00000, 0	0.00, 0	0.00
0.00000, 0	0.00, 0	0.00
0.00000, 0	0.00, 0	0.00
0.00000, 0	0.00, 0	0.00
0.00000, 0	0.00, 0	0.00

$$C_1 = 1, C_2 = 1$$

$$C_1 = 1, C_2 = 1$$

TABLE 2

See Appendix Table 2 for details.

See Appendix Table 2 for details.

$$C_1 = 1, C_2 = 1$$

β	α	γ
0.00000, 0	0.00, 0	0.00
0.00000, 0	0.00, 0	0.00
0.00000, 0	0.00, 0	0.00
0.00000, 0	0.00, 0	0.00
0.00000, 0	0.00, 0	0.00
0.00000, 0	0.00, 0	0.00

$$C_1 = 1, C_2 = 1$$

$$C_1 = 1, C_2 = 1$$

TABLE 3

See Appendix Table 3 for details.

See Appendix Table 3 for details.

$$C_1 = 1, C_2 = 1$$

β	α	γ
0.00000, 0	0.00, 0	0.00
0.00000, 0	0.00, 0	0.00
0.00000, 0	0.00, 0	0.00
0.00000, 0	0.00, 0	0.00
0.00000, 0	0.00, 0	0.00
0.00000, 0	0.00, 0	0.00

$$C_1 = 1, C_2 = 1$$

$$C_1 = 1, C_2 = 1$$

TABLE 4

See Appendix Table 4 for details.

See Appendix Table 4 for details.

$$C_1 = 1, C_2 = 1$$

β	α	γ
0.00000, 0	0.00, 0	0.00
0.00000, 0	0.00, 0	0.00
0.00000, 0	0.00, 0	0.00
0.00000, 0	0.00, 0	0.00
0.00000, 0	0.00, 0	0.00
0.00000, 0	0.00, 0	0.00

$$C_1 = 1, C_2 = 1$$

$$C_1 = 1, C_2 = 1$$

Table XXXI

0.0625 N Sodium Ethylate and
0.25 N Methyl Iodide at 0°C.

A = 10.00 B = 40.00

t	x	K _v
857	5.530	0.0002544
908	5.720	0.0002548
1031	6.160	0.0002553

Average = 0.0002548

K_n = 0.004077

Table XXXII

0.03125 N Sodium Ethylate and
0.25 N Methyl Iodide at 0°C.

A = 10.01 B = 80.00

t	x	K _v
673	4.970	0.0001321
751	5.338	0.0001319
821	5.563	0.0001287
899	5.874	0.0001284
966	6.314	0.0001353
1095	6.714	0.0001337

Average = 0.0001317

K_n = 0.004212

Table XXXIII

0.03125 N Sodium Ethylate and
0.25 N Methyl Iodide at 0°C.

A = 10.02 B = 80.00

t	x	K _v
220	2.032	0.0001305*
270	2.428	0.0001308
300	2.660	0.0001310
1327	7.180	0.0001258
1493	7.644	0.0001283

Average = 0.0001293

K_n = 0.004130

Table XXXIV

0.03125 N Sodium Ethylate and
0.125 N Methyl Iodide at 0°C.

A = 10.02 B = 40.00

t	x	K _v
1462	5.124	0.0001334
1670	5.524	0.0001306
1832	5.804	0.0001296
3020	7.476	0.0001288

Average = 0.0001304

K_n = 0.004165

* A = 10.12

1920m. 3 = 107

	2	3
2000000,0	100,0	100
2000000,0	100,0	100
2000000,0	100,0	100
2000000,0	100,0	100
2000000,0	100,0	100

$\text{C}_{60}\text{H}_8\text{O} = 96$

11. 1997年10月1日

Table XXXV

Summary of Sodium Ethylate and Methyl Iodide at 0°C.

V	K_n	Average K_n
1	0.003399 0.003364	0.003381
2	0.003400 0.003420 0.003485	0.003435
4	0.003679 0.003631	0.003650
8	0.003835 0.003854	0.003840
16	0.003919 0.003985 0.004063 0.004077	0.004011
32	0.004212 0.004130 0.004165	0.004169

Table XXXVI

Sodium Ethylate and Methyl Iodide at 0°C.

V	α	$1 - \alpha$	K_n found	K_n calculated	Per cent. error
1	0.136	0.864	0.003281	0.003282	-0.03
2	0.237	0.763	0.003435	0.003466	-0.90
4	0.335	0.665	0.003650	0.003645	+0.13
8	0.435	0.565	0.003840	0.003828	+0.31
16	0.531	0.469	0.004011	0.004004	+0.17
32	0.633	0.367	0.004169	0.004190	-0.50

Sample and description	Year	$\alpha = 1$	α	β
2011-2012	2012	0.01	0.01	1
2013-2014	2014	0.01	0.01	2
2015-2016	2016	0.01	0.01	3
2017-2018	2018	0.01	0.01	4
2019-2020	2020	0.01	0.01	5
2021-2022	2022	0.01	0.01	6

Table XXXVII

K_1 and K_m for Sodium Ethylate and Methyl Iodide at 0°C.

V	K_1	K_m
1 : 3	0.004598	0.003074
1 : 4	0.004683	0.003010
1 : 8	0.004696	0.003027
1 : 16	0.004878	0.003030
1 : 32	0.004624	0.003038
2 : 4	0.005109	0.002915
2 : 8	0.004995	0.002950
2 : 16	0.004930	0.002970
2 : 32	0.004849	0.002995
4 : 8	0.004914	0.003014
4 : 16	0.004874	0.003033
4 : 32	0.004808	0.003066
8 : 16	0.004846	0.003065
8 : 32	0.004778	0.003116
16 : 32	0.004737	0.003168

Average = 0.004861 0.003033

Potassium Ethylate and Methyl Iodide at 0°

Table XXXVIII

0.35 N Potassium Ethylate and
0.35 N Methyl Iodide at 0°

A = 10.00

t	x	K _v
110	0.972	0.000980
190	1.600	0.001002
290	2.368	0.001015
350	2.604	0.001006
390	2.804	0.000997
1430	5.936	0.001024
1590	6.196	0.001024

Average = 0.001006

K_n = 0.004024

Table XXXIX

0.35 N Potassium Ethylate and
0.35 N Methyl Iodide at 0°

A = 10.02

t	x	K _v
130	1.086	0.001014
180	1.506	0.000984
300	2.394	0.000991
440	3.026	0.000985
520	3.418	0.000997
373	2.710	0.000995
1371	5.626	0.001010

Average = 0.000997

K_n = 0.003982

Table XL

0.25 N Potassium Ethylate and
0.50 N Methyl Iodide at 0°.

A = 9.97 B = 20.00

t	x	K _v
120	2.066	0.001030
150	2.490	0.001022
180	2.870	0.001019
190	2.978	0.001012
210	3.234	0.001021

Average = 0.001019

K_n = 0.004088

Table XLI

0.1875 N Potassium Ethylate and
0.50 N Methyl Iodide at 0°.

A = 9.99 B = 26.67

t	x	K _v
90	1.639	0.0007705
130	2.279	0.0007813
160	2.679	0.0007724
180	2.940	0.0007713
220	3.436	0.0007721
260	3.890	0.0007722

Average = 0.0007735

K_n = 0.004133

K_n = 0.004133

Table XLII

0.1875 N Potassium Ethylate and
0.35 N Methyl Iodide at 0°.

A = 10.03 B = 13.33

t	x	K _v
130	1.119	0.0007755
150	1.375	0.0007818
200	1.753	0.0007769
250	3.116	0.0007790
320	3.564	0.0007755
370	3.857	0.0007728
480	3.417	0.0007630

Average = 0.0007749

K_n = 0.004134

Table XLIII

0.135 N Potassium Ethylate and
0.50 N Methyl Iodide at 0°.

A = 10.03 B = 40.00

t	x	K _v
360	4.044	0.0005278
320	4.692	0.0005292
380	5.360	0.0005304
450	5.820	0.0005293
503	6.304	0.0005288

Average = 0.0005291

K_n = 0.004234

Table XLIV

0.135 N Potassium Ethylate and
0.50 N Methyl Iodide at 0°.

A = 10.00 B = 40.00

t	x	K _v
180	3.192	0.0005285
240	3.832	0.0005311
290	4.376	0.0005282
330	4.792	0.0005299
360	5.064	0.0005280
390	5.360	0.0005331
460	5.880	0.0005271

Average = 0.0005294

K_n = 0.004235

Table XLV

0.135 N Potassium Ethylate and
0.50 N Methyl Iodide at 0°.

A = 10.00 B = 40.00

t	x	K _v
160	3.784	0.0005291
190	3.200	0.0005301
230	3.720	0.0005326
250	3.968	0.0005345
267	4.153	0.0005338
300	4.512	0.0005336

Average = 0.0005331

K_n = 0.004257

Table 1111

Calculation of the average value of the

of the average value of the

 $\sigma_{\text{avg}} = \sigma$ $\sigma_{\text{avg}} = 1$

σ	σ	σ
0.00000,0	0.00000	0.00000
0.00000,0	0.00000	0.00000
0.00000,0	0.00000	0.00000
0.00000,0	0.00000	0.00000
0.00000,0	0.00000	0.00000

Average = 0.00000

Standard = 0.00000

Table 1112

Calculation of the average value of the

of the average value of the

 $\sigma_{\text{avg}} = \sigma$ $\sigma_{\text{avg}} = 1$

σ	σ	σ
0.00000,0	0.00000	0.00000
0.00000,0	0.00000	0.00000
0.00000,0	0.00000	0.00000
0.00000,0	0.00000	0.00000
0.00000,0	0.00000	0.00000
0.00000,0	0.00000	0.00000

Average = 0.00000

Standard = 0.00000

Table 1113

Calculation of the average value of the

of the average value of the

 $\sigma_{\text{avg}} = \sigma$ $\sigma_{\text{avg}} = 1$

σ	σ	σ
0.00000,0	0.00000	0.00000
0.00000,0	0.00000	0.00000
0.00000,0	0.00000	0.00000
0.00000,0	0.00000	0.00000
0.00000,0	0.00000	0.00000
0.00000,0	0.00000	0.00000

Average = 0.00000

Standard = 0.00000

Table 1114

Calculation of the average value of the

of the average value of the

 $\sigma_{\text{avg}} = \sigma$ $\sigma_{\text{avg}} = 1$

σ	σ	σ
0.00000,0	0.00000	0.00000
0.00000,0	0.00000	0.00000
0.00000,0	0.00000	0.00000
0.00000,0	0.00000	0.00000
0.00000,0	0.00000	0.00000
0.00000,0	0.00000	0.00000

Average = 0.00000

Standard = 0.00000

Table XLVI

0.09375 N Potassium Ethylate and
0.50 N Methyl Iodide at 0°.

A = 10.00 B = 53.33

t	x	K _v
120	2.367	0.0004108
180	3.142	0.0004056
220	3.685	0.0004070
260	4.187	0.0004089
365	5.364	0.0004068
420	5.733	0.0004055

Average = 0.0004074

K_n = 0.004347

Table XLVII

0.09375 N Potassium Ethylate and
0.50 N Methyl Iodide at 0°.

A = 10.04 B = 53.33

t	x	K _v
180	3.193	0.0004135
210	3.598	0.0004130
260	4.306	0.0004109
320	4.824	0.0004058

Average = 0.0004108

K_n = 0.004362

Table XLVIII

0.09375 N Potassium Ethylate and
0.50 N Methyl Iodide at 0°.

A = 10.00 B = 53.33

t	x	K _v
120	2.356	0.0004089
180	3.131	0.0004039
220	3.664	0.0004039
370	4.393	0.0004076
420	5.723	0.0004043

Average = 0.0004057

K_n = 0.004329

Table XLIX

0.0625 N Potassium Ethylate and
0.50 N Methyl Iodide at 0°.

A = 10.00 B = 80.00

t	x	K _v
120	2.376	0.0002836
150	2.856	0.0002827
190	3.448	0.0002821
250	4.200	0.0002803
300	4.728	0.0002757
360	5.376	0.0002784
420	5.904	0.0002774

Average = 0.0002800

K_n = 0.004480

Table 11

Logarithmic correlation of $\log \rho$ and $\log \eta$

$\log \rho$ in g/cm³, $\log \eta$ in poise

$$\log \rho = 1.0 \quad \log \eta = 1.0$$

$\log \rho$	$\log \eta$	$\log \rho$
0.00000	0.00000	0.00000
0.00001	0.00001	0.00001
0.00002	0.00002	0.00002
0.00003	0.00003	0.00003
0.00004	0.00004	0.00004

$$\log \rho = 1.0 \quad \log \eta = 1.0$$

$$\log \rho = 1.0 \quad \log \eta = 1.0$$

Table 12

Logarithmic correlation of $\log \rho$ and $\log \eta$

$\log \rho$ in g/cm³, $\log \eta$ in poise

$$\log \rho = 1.0 \quad \log \eta = 1.0$$

$\log \rho$	$\log \eta$	$\log \rho$
0.00000	0.00000	0.00000
0.00001	0.00001	0.00001
0.00002	0.00002	0.00002
0.00003	0.00003	0.00003
0.00004	0.00004	0.00004
0.00005	0.00005	0.00005

$$\log \rho = 1.0 \quad \log \eta = 1.0$$

$$\log \rho = 1.0 \quad \log \eta = 1.0$$

Table 13

Logarithmic correlation of $\log \rho$ and $\log \eta$

$\log \rho$ in g/cm³, $\log \eta$ in poise

$$\log \rho = 1.0 \quad \log \eta = 1.0$$

$\log \rho$	$\log \eta$	$\log \rho$
0.00000	0.00000	0.00000
0.00001	0.00001	0.00001
0.00002	0.00002	0.00002
0.00003	0.00003	0.00003
0.00004	0.00004	0.00004
0.00005	0.00005	0.00005
0.00006	0.00006	0.00006
0.00007	0.00007	0.00007
0.00008	0.00008	0.00008
0.00009	0.00009	0.00009

$$\log \rho = 1.0 \quad \log \eta = 1.0$$

$$\log \rho = 1.0 \quad \log \eta = 1.0$$

Table 14

Logarithmic correlation of $\log \rho$ and $\log \eta$

$\log \rho$ in g/cm³, $\log \eta$ in poise

$$\log \rho = 1.0 \quad \log \eta = 1.0$$

$\log \rho$	$\log \eta$	$\log \rho$
0.00000	0.00000	0.00000
0.00001	0.00001	0.00001
0.00002	0.00002	0.00002
0.00003	0.00003	0.00003
0.00004	0.00004	0.00004
0.00005	0.00005	0.00005

$$\log \rho = 1.0 \quad \log \eta = 1.0$$

$$\log \rho = 1.0 \quad \log \eta = 1.0$$

Table L

0.0635 N Potassium Ethylate and
0.50 N Methyl Iodide at 0°.

A = 10.08 B = 80.00

t	x	K _v
180	3.272	0.0003808
330	3.938	0.0003778
390	4.672	0.0003796
350	5.304	0.0003794
430	6.008	0.0003777
500	6.544	0.0003774

Average = 0.0003783

K_n = 0.004424

Table LI

0.0635 N Potassium Ethylate and
0.1343 N Methyl Iodide at 0°.

A = 10.07 B = 21.47

t	x	K _v
835	3.638	0.0003777
910	3.838	0.0003745
935	3.894	0.0003726
1010	4.134	0.0003743
1080	4.302	0.0003736
1240	4.702	0.0003718
1365	4.936	0.0003694

Average = 0.0003733

K_n = 0.004343

Table LII

0.0635 N Potassium Ethylate and
0.35 N Methyl Iodide at 0°.

A = 10.03 B = 40.00

t	x	K _v
600	4.526	0.0003677
630	4.734	0.0003722
670	4.903	0.0003697
720	5.174	0.0003727
780	5.406	0.0003699

Average = 0.0003704

K_n = 0.004300

Table LIII

0.0635 N Potassium Ethylate and
0.50 N Methyl Iodide at 0°.

A = 10.03 B = 80.00

t	x	K _v
130	2.478	0.0003781
180	3.152	0.0003715
310	3.598	0.0003718
360	4.303	0.0003783
340	5.214	0.0003808
420	5.902	0.0003768

Average = 0.0003763

K_n = 0.004405

Abstract

$$\partial\phi_1/\partial\alpha = 0 \quad \partial\phi_2/\partial\alpha = 0$$

Table LIV

0.03135 N Potassium Ethylate and
0.135 N Methyl Iodide at 0°.

A = 10.03 B = 40.00

t	x	K _v
890	3.764	0.0001397
1000	4.093	0.0001393
1110	4.444	0.0001409
1200	4.604	0.0001373
1330	4.990	0.0001403
1413	5.353	0.0001417

Average = 0.0001398

K_n = 0.004466

Table LV

0.03135 N Potassium Ethylate and
0.35 N Methyl Iodide at 0°.

A = 10.00 B = 80.00

t	x	K _v
650	4.936	0.0001355
700	5.264	0.0001385
730	5.456	0.0001405
760	5.534	0.0001400
800	5.713	0.0001379
840	5.888	0.0001381
990	6.424	0.0001363

Average = 0.0001381

K_n = 0.004419

Table LVI

0.03135 N Potassium Ethylate and
0.35 N Methyl Iodide at 0°.

A = 10.03 B = 80.00

t	x	K _v
180	1.830	0.0001430
240	2.350	0.0001416
330	2.894	0.0001360
360	3.330	0.0001383
430	3.743	0.0001430
480	4.094	0.0001409

Average = 0.0001403

K_n = 0.004476

Table LVII

0.03135 N Potassium Ethylate and
0.35 N Methyl Iodide at 0°.

A = 10.01 B = 80.00

t	x	K _v
160	1.643	0.0001415
230	2.358	0.0001475
370	2.554	0.0001390
340	3.130	0.0001409
390	3.554	0.0001441
450	3.933	0.0001420

Average = 0.0001425

K_n = 0.004556

TABLE

See explanatory memorandum to Table 1

1. For values of x and y

$$x, y = 1 \quad x, y = 2$$

x^2	y	z
1.000000	1.000000	1.000000
1.000000	1.000000	1.000000
1.000000	1.000000	1.000000
1.000000	1.000000	1.000000
1.000000	1.000000	1.000000
1.000000	1.000000	1.000000
1.000000	1.000000	1.000000

$$1.000000 = 1.000000$$

$$1.000000 = 1.000000$$

TABLE

See explanatory memorandum to Table 1

1. For values of x and y

$$x, y = 1 \quad x, y = 2$$

x^2	y	z
1.000000	1.000000	1.000000
1.000000	1.000000	1.000000
1.000000	1.000000	1.000000
1.000000	1.000000	1.000000
1.000000	1.000000	1.000000
1.000000	1.000000	1.000000
1.000000	1.000000	1.000000

$$1.000000 = 1.000000$$

$$1.000000 = 1.000000$$

TABLE

See explanatory memorandum to Table 1

TABLE

See explanatory memorandum to Table 1

1. For values of x and y

$$x, y = 1 \quad x, y = 2$$

x^2	y	z
1.000000	1.000000	1.000000
1.000000	1.000000	1.000000
1.000000	1.000000	1.000000
1.000000	1.000000	1.000000
1.000000	1.000000	1.000000
1.000000	1.000000	1.000000
1.000000	1.000000	1.000000

$$1.000000 = 1.000000$$

$$1.000000 = 1.000000$$

TABLE

See explanatory memorandum to Table 1

1. For values of x and y

$$x, y = 1 \quad x, y = 2$$

x^2	y	z
1.000000	1.000000	1.000000
1.000000	1.000000	1.000000
1.000000	1.000000	1.000000
1.000000	1.000000	1.000000
1.000000	1.000000	1.000000
1.000000	1.000000	1.000000
1.000000	1.000000	1.000000

$$1.000000 = 1.000000$$

$$1.000000 = 1.000000$$

Table LVIII

0.03135 N Potassium Ethylate and
0.35 N Methyl Iodide at 0°.

A = 10.00 B = 80.00

t	x	K _v
140	1.448	0.0001409
200	1.952	0.0001374
250	2.434	0.0001410
320	2.960	0.0001398
380	3.440	0.0001419
430	3.736	0.0001394

Average = 0.0001401

K_n = 0.004483

Table LIX

0.03135 N Potassium Ethylate and
0.50 N Methyl Iodide at 0°.

A = 10.00 B = 160.00

t	x	K _v
190	3.528	0.0001446
220	3.968	0.0001454
270	4.568	0.0001433
360	5.480	0.0001404
430	6.080	0.0001423

Average = 0.0001432

K_n = 0.004582

Table LX

Summary of Potassium Ethylate and Methyl Iodide at 0°.

V	K _n	Average K _n
4	0.004034 0.003983 0.004088	0.004031
16/3	0.004133 0.004134 0	0.004134
8	0.004234 0.004235 0.004258	0.004239
32/3	0.004347 0.004362 0.004329	0.004346
16	0.004480 0.004300 0.004434 0.004343 0.004405	0.004390
32	0.004466 0.004419 0.004582 0.004476 0.004556 0.004482	0.004497

TABLE 1

TABLE 2

and the other two are the same as in Table 1

and the other two are the same as in Table 1

10.0 to 10.0, 10.0 to 10.0

10.0 to 10.0, 10.0 to 10.0

10.0 to 10.0, 10.0 to 10.0

10.0 to 10.0, 10.0 to 10.0

10.0 to 10.0, 10.0 to 10.0

10.0 to 10.0, 10.0 to 10.0

10.0 to 10.0, 10.0 to 10.0

10.0 to 10.0, 10.0 to 10.0

10.0 to 10.0, 10.0 to 10.0

10.0 to 10.0, 10.0 to 10.0

10.0 to 10.0, 10.0 to 10.0

10.0 to 10.0, 10.0 to 10.0

10.0 to 10.0, 10.0 to 10.0

10.0 to 10.0, 10.0 to 10.0

10.0 to 10.0, 10.0 to 10.0

10.0 to 10.0, 10.0 to 10.0

10.0 to 10.0, 10.0 to 10.0

10.0 to 10.0, 10.0 to 10.0

TABLE 3

10.0 to 10.0, 10.0 to 10.0, 10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

10.0 to 10.0

Table LXI

Potassium Ethylate and Methyl Iodide at 0°C.

V	α	$1 - \alpha$	K_n found	K_n calculated	Per cent. error
4	0.365	0.635	0.004031	0.004055	-0.63
16/3	0.401	0.599	0.004124	0.004123	+0.02
8	0.459	0.541	0.004339	0.004333	+0.12
33/3	0.491	0.509	0.004346	0.004293	+1.26
16	0.546	0.454	0.004390	0.004397	-0.10
33	0.637	0.363	0.004497	0.004568	-1.55

Table LXII

K_1 and K_m for Potassium Ethylate and Methyl Iodide at 0°.

V	K_1	K_m
4 : 8	0.00543	0.00320
4 : 33/3	0.00562	0.00310
4 : 16	0.00528	0.00329
4 : 33	0.00513	0.00340
16/3 : 33/3	0.00564	0.00306
16/3 : 16	0.00522	0.00335
16/3 : 33	0.00508	0.00346
8 : 16	0.00517	0.00345
8 : 33	0.00502	0.00359
16 : 33	0.00493	0.00377

Average = 0.00525 0.00337

Sodium Ethylate and Methyl Iodide at 25°C.

Table LXIII

0.5 N Sodium Ethylate and
0.5 N Methyl Iodide at 25°C.

A = 10.00

t	x	K_v
20	4.343	0.03839
30	5.337	0.03815
40	5.999	0.03745
50	6.495	0.03714
60	6.901	0.03711
70	7.187	0.03650

Average = 0.03746

 $K_n = 0.07493$

Table LXIV

0.5 N Sodium Ethylate and
0.5095 N Methyl Iodide at 25°C.

A = 10.00 B = 10.19

t	x	K_v
15	3.657	0.03715
24	4.830	0.03786
33	5.600	0.03745
43	6.166	0.03691
51	6.586	0.03634
69	7.307	0.03564
97	7.815	0.03500

Average = 0.03662

 $K_n = 0.07324$

Table LXV

0.5 N Sodium Ethylate and
0.35 N Methyl Iodide at 25°C.

A = 20.00 B = 10.00

t	x	K_v
15	4.044	0.03894
25	5.492	0.03805
35	6.476	0.03723
55	7.818	0.03722
65	8.196	0.03646
75	8.520	0.03613

Average = 0.03735

 $K_n = 0.07471$

Table LXVI

0.5 N Sodium Ethylate and
0.5 N Methyl Iodide at 25°C.

A = 9.989

t	x	K_v
10	2.891	0.04073
30	4.375	0.03896
30	5.367	0.03870
40	6.027	0.03803
50	6.527	0.03770
60	6.889	0.03703
70	7.175	0.03643

Average = 0.03832

 $K_n = 0.07652$

1000

STAT. 0 = 1

Table LXVII

0.5 N Sodium Ethylate and
0.35 N Methyl Iodide at 25°C.

A = 20.04 B = 10.00

t	x	K_v
20	4.780	0.03768
30	6.032	0.03766
37	6.876	0.03755
43	7.120	0.03740
50	7.520	0.03687
59	7.932	0.03626
68	8.308	0.03643

Average = 0.03712

$K_n = 0.07409$

Table LXVIII

0.35 N Sodium Ethylate and
0.35 N Methyl Iodide at 25°C.

A = 10.00

t	x	K_v
10	1.685	0.03026
20	2.909	0.03051
30	3.760	0.03008
40	4.446	0.03001
50	5.000	0.03000
60	5.430	0.01980
70	5.793	0.01967

Average = 0.03005

$K_n = 0.08020$

Table LXIX

0.35 N Sodium Ethylate and
0.50 N Methyl Iodide at 25°C.

A = 10.02 B = 20.00

t	x	K_v
15	4.152	0.02024
25	5.668	0.02009
35	6.682	0.01988
45	7.418	0.01972
55	7.956	0.01934
65	8.362	0.01942
75	8.698	0.01947

Average = 0.01974

$K_n = 0.07880$

Table LXX

0.25 N Sodium Ethylate and
0.25 N Methyl Iodide at 25°C.

A = 9.956

t	x	K_v
20	2.868	0.02023
25	3.340	0.02019
30	3.758	0.02021
40	4.453	0.02023
50	4.980	0.02002
60	5.420	0.01991

Average = 0.02013

$K_n = 0.08086$

TABLE 1

For calculation of the χ^2 value
of the χ^2 test for the χ^2 test
of the χ^2 test for the χ^2 test

χ^2	χ^2	χ^2
0.0000	0.0000	0.0000
0.0001	0.0001	0.0001
0.0002	0.0002	0.0002
0.0003	0.0003	0.0003
0.0004	0.0004	0.0004
0.0005	0.0005	0.0005
0.0006	0.0006	0.0006
0.0007	0.0007	0.0007

0.0008 = 0.0008

0.0009 = 0.0009

TABLE 2

For calculation of the χ^2 value
of the χ^2 test for the χ^2 test
of the χ^2 test for the χ^2 test

χ^2	χ^2	χ^2
0.0000	0.0000	0.0000
0.0001	0.0001	0.0001
0.0002	0.0002	0.0002
0.0003	0.0003	0.0003
0.0004	0.0004	0.0004
0.0005	0.0005	0.0005
0.0006	0.0006	0.0006
0.0007	0.0007	0.0007

0.0008 = 0.0008

0.0009 = 0.0009

TABLE 3

For calculation of the χ^2 value
of the χ^2 test for the χ^2 test
of the χ^2 test for the χ^2 test

χ^2	χ^2	χ^2
0.0000	0.0000	0.0000
0.0001	0.0001	0.0001
0.0002	0.0002	0.0002
0.0003	0.0003	0.0003
0.0004	0.0004	0.0004
0.0005	0.0005	0.0005
0.0006	0.0006	0.0006
0.0007	0.0007	0.0007

0.0008 = 0.0008

0.0009 = 0.0009

TABLE 4

For calculation of the χ^2 value
of the χ^2 test for the χ^2 test
of the χ^2 test for the χ^2 test

χ^2	χ^2	χ^2
0.0000	0.0000	0.0000
0.0001	0.0001	0.0001
0.0002	0.0002	0.0002
0.0003	0.0003	0.0003
0.0004	0.0004	0.0004
0.0005	0.0005	0.0005
0.0006	0.0006	0.0006
0.0007	0.0007	0.0007

0.0008 = 0.0008

0.0009 = 0.0009

Table LXXI

0.35 N Sodium Ethylate and
0.35 N Methyl Iodide at 25°C.

A = 10.01

t	x	K_v
20	2.918	0.02057
35	3.354	0.02015
30	3.768	0.02012
40	4.455	0.02005
46	4.795	0.01998
52	5.076	0.01978

Average = 0.02011

$K_n = 0.08035$

Table LXXII

0.125 N Sodium Ethylate and
0.125 N Methyl Iodide at 25°C.

A = 10.00

t	x	K_v
40	2.959	0.01051
55	3.691	0.01064
70	4.223	0.01045
80	4.575	0.01054
90	4.852	0.01047
100	5.111	0.01045

Average = 0.01051

$K_n = 0.08406$

Table LXXIII

0.125 N Sodium Ethylate and
0.25 N Methyl Iodide at 25°C.

A = 10.00 B = 20.00

t	x	K_v
15	2.594	0.01075
25	3.836	0.01082
35	4.830	0.01094
45	5.548	0.01075
55	6.163	0.01072
65	6.686	0.01072
75	7.103	0.01067

Average = 0.01077

$K_n = 0.08616$

Table LXXIV

0.125 N Sodium Ethylate and
0.25 N Methyl Iodide at 25°C.

A = 10.03 B = 20.00

t	x	K_v
20	3.318	0.01107
26	3.922	0.01098
34	4.734	0.01090
42	5.382	0.01091
47	5.734	0.01091
57	6.318	0.01084

Average = 0.01094

$K_n = 0.08722$

TABLE 111

See analysis number 7 111.0
 111.0 is subject to 111.0

 $111.0 = 111.0$

χ^2	α	β
111.0, 0	111.0	111.0
111.0, 0	111.0	111.0
111.0, 0	111.0	111.0
111.0, 0	111.0	111.0
111.0, 0	111.0	111.0
111.0, 0	111.0	111.0
111.0, 0	111.0	111.0

 $111.0 = 111.0$
 $111.0 = 111.0$

TABLE 112

See analysis number 7 112.0
 112.0 is subject to 112.0

 $112.0 = 112.0$

χ^2	α	β
112.0, 0	112.0	112.0
112.0, 0	112.0	112.0
112.0, 0	112.0	112.0
112.0, 0	112.0	112.0
112.0, 0	112.0	112.0
112.0, 0	112.0	112.0
112.0, 0	112.0	112.0

 $112.0 = 112.0$
 $112.0 = 112.0$
 $112.0 = 112.0$

TABLE 113

See analysis number 7 113.0
 113.0 is subject to 113.0

 $113.0 = 113.0$

χ^2	α	β
113.0, 0	113.0	113.0
113.0, 0	113.0	113.0
113.0, 0	113.0	113.0
113.0, 0	113.0	113.0
113.0, 0	113.0	113.0
113.0, 0	113.0	113.0
113.0, 0	113.0	113.0

 $113.0 = 113.0$
 $113.0 = 113.0$

TABLE 114

See analysis number 7 114.0
 114.0 is subject to 114.0

 $114.0 = 114.0$

χ^2	α	β
114.0, 0	114.0	114.0
114.0, 0	114.0	114.0
114.0, 0	114.0	114.0
114.0, 0	114.0	114.0
114.0, 0	114.0	114.0
114.0, 0	114.0	114.0
114.0, 0	114.0	114.0

 $114.0 = 114.0$
 $114.0 = 114.0$

Table LXXV

0.125 N Sodium Ethylate and
0.35 N Methyl Iodide at 25°C.

A = 10.00 B = 20.00

t	x	K _v
15	2.648	0.01103
25	3.848	0.01088
33	4.560	0.01094
38	5.080	0.01095
46	5.656	0.01090
52	5.992	0.01073
58	6.312	0.01066

Average = 0.01087

K_n = 0.08696

Table LXXVI

0.0635 N Sodium Ethylate and
0.25 N Methyl Iodide at 25°C.

A = 10.00 B = 41.00

t	x	K _v
15	2.889	0.005475
25	4.311	0.005842
35	5.366	0.005792
45	6.180	0.005725
55	6.884	0.005759
65	7.435	0.005934

Average = 0.005754

K_n = 0.09206

Table LXXVII

0.0625 N Sodium Ethylate and
0.25 N Methyl Iodide at 25°C.

A = 10.00 B = 40.00

t	x	K _v
15	2.800	0.005678
25	4.181	0.005746
35	5.256	0.005771
45	6.105	0.005768
55	7.000	0.006122
65	7.436	0.005911
75	7.724	0.005622

Average = 0.005802

K_n = 0.09198

Table LXXVIII

0.03125 N Sodium Ethylate and
0.25 N Methyl Iodide at 25°C.

A = 9.912 B = 80.00

t	x	K _v
15	3.001	0.003029
25	4.550	0.003144
35	5.607	0.003075
45	6.305	0.003088
55	7.184	0.003074
65	7.741	0.003081

Average = 0.003082

K_n = 0.09948

TABLE 1

Low frequency modes (X 1000)

Low frequency modes (X 1000)

0.000000 0.000000

μ	λ	ν
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000

0.000000 = 0.000000

0.000000 = 0.000000

TABLE 2

Low frequency modes (X 1000)

Low frequency modes (X 1000)

0.000000 0.000000

μ	λ	ν
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000

0.000000 = 0.000000

0.000000 = 0.000000

TABLE 3

Low frequency modes (X 1000)

Low frequency modes (X 1000)

0.000000 0.000000

μ	λ	ν
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000

0.000000 = 0.000000

0.000000 = 0.000000

TABLE 4

Low frequency modes (X 1000)

Low frequency modes (X 1000)

0.000000 0.000000

μ	λ	ν
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000
0.000000	0.000000	0.000000

0.000000 = 0.000000

0.000000 = 0.000000

Table LXXIX

0.03125 N Sodium Ethylate and
0.135 N Methyl Iodide at 25°C.

A = 10.07 B = 40.00

t	x	K _v
55	4.876	0.003252
80	5.893	0.003029
100	6.783	0.003143
135	7.733	0.003104
150	8.038	0.003085
180	8.482	0.003007

Average = 0.003103

K_n = 0.09860

Table LYXX

0.03125 N Sodium Ethylate and
0.25 N Methyl Iodide at 25°C.

A = 10.00 B = 80.00

t	x	K _v
20	3.780	0.003045
25	4.467	0.003053
30	5.168	0.003144
35	5.504	0.003972
40	6.120	0.003098
45	6.595	0.003145
50	6.956	0.003137

Average = 0.003085

K_n = 0.09872

Table LXXXI

Summary of Sodium Ethylate and Methyl Iodide at 25°C.

V	V	K _n	Average K _n
	2	0.07492 0.07324 0.07471 0.07652 0.07409	0.07470
	4	0.08030 0.07880 0.08086 0.08035	0.08005
	8	0.08406 0.08616 0.08722 0.08696	0.08610
	16	0.09206 0.09198	0.09202
	32	0.09872 0.09948 0.09860	0.09893

TYPE 1000

USE KEYBOARD METHOD TO ENTER

USE AS MANY DIGITS AS DESIRED

00.00 = 0 00.00 = 0

00.00 = 0 00.00 = 0

00.00 = 0 00.00 = 0

00.00 = 0 00.00 = 0

00.00 = 0 00.00 = 0

00.00 = 0 00.00 = 0

00.00 = 0 00.00 = 0

00.00 = 0 00.00 = 0

00.00 = 0 00.00 = 0

00.00 = 0 00.00 = 0

00.00 = 0 00.00 = 0

TYPE 1000

USE KEYBOARD METHOD TO ENTER

USE AS MANY DIGITS AS DESIRED

00.00 = 0 00.00 = 0

00.00 = 0 00.00 = 0

00.00 = 0 00.00 = 0

00.00 = 0 00.00 = 0

00.00 = 0 00.00 = 0

00.00 = 0 00.00 = 0

00.00 = 0 00.00 = 0

00.00 = 0 00.00 = 0

00.00 = 0 00.00 = 0

00.00 = 0 00.00 = 0

00.00 = 0 00.00 = 0

TYPE 1000

USE AS MANY DIGITS AS DESIRED

00.00 = 0

00.00 = 0

00.00 = 0

00.00 = 0

00.00 = 0

00.00 = 0

00.00 = 0

00.00 = 0

00.00 = 0

00.00 = 0

00.00 = 0

00.00 = 0

00.00 = 0

00.00 = 0

00.00 = 0

00.00 = 0

00.00 = 0

00.00 = 0

00.00 = 0

00.00 = 0

Table LXXII

Sodium Ethylate and Methyl Iodide at 25°C.

V	α	$1 - \alpha$	K_n found	K_n calculated	Per cent. error
2	0.234	0.766	0.07470	0.07466	+0.053
4	0.312	0.688	0.08005	0.08026	-0.26
8	0.393	0.607	0.08610	0.08590	+0.23
16	0.481	0.519	0.09203	0.09191	+0.12
32	0.584	0.416	0.09893	0.09909	-0.16

Table LXXXIII

K_1 and K_m for Sodium Ethylate and Methyl Iodide at 25°C.

V	K_1	K_m
2 : 4	0.1272	0.05865
2 : 8	0.1296	0.05792
2 : 16	0.1284	0.05829
2 : 32	0.1277	0.05850
4 : 8	0.1313	0.05674
4 : 16	0.1288	0.05795
4 : 32	0.1278	0.05839
8 : 16	0.1269	0.05966
8 : 32	0.1269	0.05970
16 : 32	0.1268	0.05994

Average = 0.1281

0.05857

TABLE 1

TABLE 1. SUMMARY OF DATA FOR THE FIRST TWO YEARS

Area	Year	Area	Year	Area	Year
1.0	1960	2.0	1961	3.0	1962
4.0	1963	5.0	1964	6.0	1965
7.0	1966	8.0	1967	9.0	1968
10.0	1969	11.0	1970	12.0	1971
13.0	1972	14.0	1973	15.0	1974

TABLE 2

TABLE 2. SUMMARY OF DATA FOR THE LAST TWO YEARS

Area	Year	Area	Year
16.0	1975	17.0	1976
18.0	1977	19.0	1978
20.0	1979	21.0	1980
22.0	1981	23.0	1982
24.0	1983	25.0	1984
26.0	1985	27.0	1986
28.0	1987	29.0	1988
30.0	1989	31.0	1990
32.0	1991	33.0	1992
34.0	1993	35.0	1994

SALT EFFECT

Sodium Ethylate, Methyl Iodide and Sodium Iodide at 25 °C.

Table LXXXIV

0.35 N Sodium Ethylate,
0.35 N Methyl Iodide and
0.35 N Sodium Iodide at 25°C.

$$A = 9.971$$

t	x	K_v
15	3.067	0.01743
35	3.031	0.01747
31	3.500	0.01744
40	4.097	0.01744
48	4.538	0.01733
57	4.953	0.01733
66	5.307	0.01734

$$\text{Average} = 0.01738$$

$$K_n = 0.06972$$

Table LXXXV

0.35 N Sodium Ethylate,
0.35 N Methyl Iodide and
0.35 N Sodium Iodide at 25°C.

$$A = 10.01$$

t	x	K_v
15	3.078	0.01747
33	3.874	0.01751
31	3.506	0.01739
41	4.170	0.01743
48	4.547	0.01734
56	4.930	0.01734
64	5.352	0.01733

$$\text{Average} = 0.01740$$

$$K_n = 0.06953$$

Table LXXXVI

0.35 N Sodium Ethylate,
0.35 N Methyl Iodide and
0.75 N Sodium Iodide.

$$A = 9.98$$

t	x	K_v
15	1.828	0.01495
33	3.538	0.01483
31	3.144	0.01484
41	3.750	0.01468
50	4.333	0.01473
60	4.657	0.01458
71	5.098	0.01471

$$\text{Average} = 0.01476$$

$$K_n = 0.05915$$

Table LXXXVII

0.35 N Sodium Ethylate and
0.35 N Methyl Iodide and
0.75 N Sodium Iodide.

$$A = 9.98$$

t	x	K_v
15	1.864	0.01531
34	3.638	0.01489
34	3.344	0.01483
43	3.864	0.01469
53	4.373	0.01471
63	4.804	0.01473
73	5.130	0.01463

$$\text{Average} = 0.01483$$

$$K_n = 0.05942$$

TABLE 1. Results of the analysis of variance for the different treatments.

Treatment

Treatment

1. Control (Control)

1. Control (Control)

2. 100% N (100% N)

2. 100% N (100% N)

3. 100% P (100% P)

3. 100% P (100% P)

100% N = 1

100% P = 1

Treatment	1	2
Control	100	100
100% N	100	100
100% P	100	100
100% N + 100% P	100	100
100% N + 100% P + 100% K	100	100
100% N + 100% P + 100% K + 100% S	100	100
100% N + 100% P + 100% K + 100% S + 100% Z	100	100

100% N = 100%

100% P = 100%

Treatment	1	2
Control	100	100
100% N	100	100
100% P	100	100
100% N + 100% P	100	100
100% N + 100% P + 100% K	100	100
100% N + 100% P + 100% K + 100% S	100	100
100% N + 100% P + 100% K + 100% S + 100% Z	100	100

100% N = 100%

100% P = 100%

Treatment

Treatment

1. Control (Control)

1. Control (Control)

2. 100% N (100% N)

2. 100% N (100% N)

3. 100% P (100% P)

3. 100% P (100% P)

100% N = 1

100% P = 1

Treatment	1	2
Control	100	100
100% N	100	100
100% P	100	100
100% N + 100% P	100	100
100% N + 100% P + 100% K	100	100
100% N + 100% P + 100% K + 100% S	100	100
100% N + 100% P + 100% K + 100% S + 100% Z	100	100

100% N = 100%

100% P = 100%

Treatment	1	2
Control	100	100
100% N	100	100
100% P	100	100
100% N + 100% P	100	100
100% N + 100% P + 100% K	100	100
100% N + 100% P + 100% K + 100% S	100	100
100% N + 100% P + 100% K + 100% S + 100% Z	100	100

100% N = 100%

100% P = 100%

Table LXXXV II

0.135 N Sodium Ethylate,
0.35 N Methyl Iodide and
0.135 N Sodium Iodide.

A = 9.98 B = 20.00

t	x	K_v
15	2.394	0.009351
20	3.036	0.009863
38	3.778	0.009473
34	4.337	0.009540
42	4.930	0.009409
50	5.484	0.009498
58	5.916	0.009404

Average = 0.009491

$K_n = 0.07608$

Table LXXXVI

0.135 N Sodium Ethylate,
0.25 N Methyl Iodide and
0.125 N Sodium Iodide.

A = 9.976 B = 20.00

t	x	K_v
15	2.326	0.009408
23	3.188	0.009580
39	3.906	0.009591
37	4.605	0.009613
44	5.121	0.009596
52	5.634	0.009555
59	5.988	0.009461

Average = 0.009541

$K_n = 0.07651$

Table XC

0.125 N Sodium Ethylate,
0.35 N Methyl Iodide and
0.375 N Sodium Iodide.

A = 9.99 B = 20.00

t	x	K_v
15	2.095	0.008395
23	2.951	0.008363
33	3.785	0.008307
39	4.356	0.008369
47	4.905	0.008362
54	5.311	0.008312
60	5.655	0.008356

Average = 0.008323

$K_n = 0.06664$

Table XCI

0.125 N Sodium Ethylate,
0.35 N Methyl Iodide and
0.375 N Sodium Iodide.

A = 10.01 B = 20.00

t	x	K_v
15	2.103	0.008325
23	2.978	0.008358
30	3.656	0.008433
38	4.287	0.008380
47	4.899	0.008340
55	5.390	0.008362
63	5.818	0.008374

Average = 0.008367

$K_n = 0.06689$

Table XCII

0.135 N Sodium Ethylate,
0.25 N Methyl Iodide and
0.875 N Sodium Iodide.

$$A = 9.959 \quad B = 30.00$$

t	x	K_v
15	1.879	0.007299
23	2.634	0.007154
33	3.499	0.007227
40	3.993	0.007181
49	4.556	0.007142
57	4.999	0.007122
64	5.351	0.007116

$$\text{Average} = 0.007177$$

$$K_n = 0.05765$$

Table XCIII

0.135 N Sodium Ethylate,
0.25 N Methyl Iodide and
0.875 N Sodium Iodide.

$$A = 10.01 \quad B = 30.00$$

t	x	K_v
15	1.818	0.007023
25	2.774	0.007023
35	3.590	0.007051
44	4.250	0.007144
54	4.818	0.007066
63	5.290	0.007070
72	5.686	0.007025

$$\text{Average} = 0.007057$$

$$K_n = 0.05640$$

Table XCIV

0.0625 N Sodium Ethylate,
0.25 N Methyl Iodide and
0.0625 N Sodium Iodide.

$$A = 10.10 \quad B = 40.00$$

t	x	K_v
15	2.626	0.005250
23	3.550	0.005230
32	4.342	0.005218
34	4.886	0.005273
40	5.425	0.005265
46	5.852	0.005197
53	6.276	0.005199

$$\text{Average} = 0.005232$$

$$K_n = 0.08287$$

Table XCV

0.0625 N Sodium Ethylate,
0.25 N Methyl Iodide and
0.0625 N Sodium Iodide.

$$A = 10.03 \quad B = 40.00$$

t	x	K_v
15	2.604	0.005203
21	3.450	0.005279
27	4.156	0.005271
33	4.736	0.005201
38	5.214	0.005230
44	5.694	0.005209
51	6.150	0.005135

$$\text{Average} = 0.005218$$

$$K_n = 0.08232$$

(11) $\alpha = 0$

calculated values γ (100, 1

100, 1000) versus β (0.0, 0

calculated values γ (100, 1

0.0, 0.1, 0.2, 0.3, 0.4, 0.5

β	γ	α
0.0	100.0	0.0
0.1	100.0	0.1
0.2	100.0	0.2
0.3	100.0	0.3
0.4	100.0	0.4
0.5	100.0	0.5

calculated values γ (100, 1

calculated values γ (100, 1

(12) $\alpha = 0$

calculated values γ (100, 1

100, 1000) versus β (0.0, 0

calculated values γ (100, 1

0.0, 0.1, 0.2, 0.3, 0.4, 0.5

β	γ	α
0.0	100.0	0.0
0.1	100.0	0.1
0.2	100.0	0.2
0.3	100.0	0.3
0.4	100.0	0.4
0.5	100.0	0.5

calculated values γ (100, 1

calculated values γ (100, 1

(13) $\alpha = 0$

calculated values γ (100, 1

100, 1000) versus β (0.0, 0

calculated values γ (100, 1

0.0, 0.1, 0.2, 0.3, 0.4, 0.5

β	γ	α
0.0	100.0	0.0
0.1	100.0	0.1
0.2	100.0	0.2
0.3	100.0	0.3
0.4	100.0	0.4
0.5	100.0	0.5

calculated values γ (100, 1

calculated values γ (100, 1

(14) $\alpha = 0$

calculated values γ (100, 1

100, 1000) versus β (0.0, 0

calculated values γ (100, 1

0.0, 0.1, 0.2, 0.3, 0.4, 0.5

β	γ	α
0.0	100.0	0.0
0.1	100.0	0.1
0.2	100.0	0.2
0.3	100.0	0.3
0.4	100.0	0.4
0.5	100.0	0.5

calculated values γ (100, 1

calculated values γ (100, 1

Table XCVI

0.0635 N Sodium Ethylate,
0.25 N Methyl Iodide and
0.0635 N Sodium Iodide.

A = 10.03 B = 40.00

t	x	K_v
10	1.893	0.005357
20	3.305	0.005326
30	4.436	0.005162
40	5.373	0.005188
50	6.106	0.005150
60	6.731	0.005153
70	7.319	0.005107

Average = 0.005192

$K_n = 0.08381$

Table XCVII

0.0635 N Sodium Ethylate,
0.25 N Methyl Iodide and
0.1875 N Sodium Iodide.

A = 10.08 B = 40.00

t	x	K_v
15	2.336	0.004568
22	3.200	0.004570
38	3.856	0.004580
34	4.468	0.004628
41	5.040	0.004587
49	5.640	0.004590
58	6.184	0.004545

Average = 0.004581

$K_n = 0.07270$

Table XCVIII

0.0635 N Sodium Ethylate,
0.25 N Methyl Iodide and
0.1875 N Sodium Iodide.

A = 10.00 B = 40.00

t	x	K_v
15	2.380	0.004444
23	3.184	0.004548
38	3.833	0.004552
34	4.438	0.004581
41	5.024	0.004581
49	5.600	0.004557
57	6.144	0.004595

Average = 0.004551

$K_n = 0.07382$

Table XCIX

0.0635 N Sodium Ethylate,
0.25 N Methyl Iodide and
0.4375 N Sodium Iodide.

A = 10.04 B = 40.00

t	x	K_v
15	2.160	0.004171
21	2.848	0.004144
38	3.576	0.004148
35	4.300	0.004125
43	4.784	0.004146
49	5.348	0.004095
56	5.660	0.004074

Average = 0.004139

$K_n = 0.06582$

Table C

0.0625 N Sodium Ethylate,
0.25 N Methyl Iodide and
0.4375 N Sodium Iodide.

A = 10.04 B = 40.00

t	x	K_v
15	2.152	0.004152
23	2.944	0.004120
39	3.659	0.004138
36	4.284	0.004133
43	4.882	0.004175
50	5.360	0.004150
56	5.720	0.004123

Average = 0.004138

$K_n = 0.06596$

Table CI

0.0625 N Sodium Ethylate,
0.25 N Methyl Iodide and
0.9375 N Sodium Iodide.

A = 10.00 B = 40.00

t	x	K_v
15	1.720	0.003322
35	2.766	0.003361
34	3.592	0.003439
44	4.357	0.003460
53	4.897	0.003409
62	5.380	0.003374

Average = 0.003394

$K_n = 0.05430$

Table CII

0.0625 N Sodium Ethylate,
0.25 N Methyl Iodide and
0.9375 N Sodium Iodide.

A = 9.99 B = 40.00

t	x	K_v
15	1.826	0.003442
24	2.738	0.003458
33	3.550	0.003490
41	4.186	0.003511
50	4.786	0.003492
59	5.290	0.003459
68	5.770	0.003455

Average = 0.003472

$K_n = 0.05561$

Table CIII

0.03125 N Sodium Ethylate,
0.25 N Methyl Iodide and
0.03125 N Sodium Iodide.

A = 10.00 B = 80.00

t	x	K_v
15	2.840	0.003836
25	4.213	0.003817
35	5.320	0.003819
45	6.226	0.003836
55	6.932	0.003834
65	7.456	0.003794
75	7.859	0.003739

Average = 0.003811

$K_n = 0.08995$

Table CIV

0.03125 N Sodium Ethylate,
0.25 N Methyl Iodide and
0.03125 N Sodium Iodide.

A = 10.00 B = 80.00

t	x	K_v
15	3.880	0.002886
25	4.336	0.002836
35	5.360	0.002852
45	6.162	0.002786
55	6.904	0.002811
65	7.432	0.002774
75	7.859	0.002740

Average = 0.002813

K_n = 0.09002

Table CV

0.03125 N Sodium Ethylate,
0.25 N Methyl Iodide and
0.09375 N Sodium Iodide.

A = 10.00 B = 80.00

t	x	K_v
30	3.160	0.002435
30	4.340	0.002445
40	5.389	0.002444
50	6.041	0.002433
60	6.709	0.002436
70	7.246	0.002439

Average = 0.002436

K_n = 0.07795

Table CVI

0.03125 N Sodium Ethylate,
0.25 N Methyl Iodide and
0.09375 N Sodium Iodide.

A = 10.00 B = 80.00

t	x	K_v
10	1.763	0.002453
20	3.205	0.002468
30	4.459	0.002438
40	5.454	0.002554
50	6.155	0.002502
60	6.814	0.002510
70	7.330	0.002499

Average = 0.002489

K_n = 0.07964

Table CVII

0.03125 N Sodium Ethylate,
0.25 N Methyl Iodide and
0.09375 N Sodium Iodide.

A = 10.07 B = 80.00

t	x	K_v
15	3.518	0.002455
22	3.470	0.002474
28	4.134	0.002444
35	4.870	0.002460
41	5.374	0.002434
49	5.974	0.002415

Average = 0.002447

K_n = 0.07776

TABLE 1

TABLE 1. Results of the first experiment.

TABLE 1. Results of the first experiment.

TABLE 1. Results of the first experiment.

TABLE 1. Results of the first experiment.

β	α	γ
0.0000	0.0000	0.0000
0.0000	0.0000	0.0000
0.0000	0.0000	0.0000
0.0000	0.0000	0.0000
0.0000	0.0000	0.0000
0.0000	0.0000	0.0000

TABLE 1. Results of the first experiment.

TABLE 1. Results of the first experiment.

TABLE 2

TABLE 2. Results of the second experiment.

TABLE 2. Results of the second experiment.

TABLE 2. Results of the second experiment.

TABLE 2. Results of the second experiment.

β	α	γ
0.0000	0.0000	0.0000
0.0000	0.0000	0.0000
0.0000	0.0000	0.0000
0.0000	0.0000	0.0000
0.0000	0.0000	0.0000
0.0000	0.0000	0.0000

TABLE 2. Results of the second experiment.

TABLE 2. Results of the second experiment.

TABLE 3

TABLE 3. Results of the third experiment.

TABLE 3. Results of the third experiment.

TABLE 3. Results of the third experiment.

TABLE 3. Results of the third experiment.

β	α	γ
0.0000	0.0000	0.0000
0.0000	0.0000	0.0000
0.0000	0.0000	0.0000
0.0000	0.0000	0.0000
0.0000	0.0000	0.0000
0.0000	0.0000	0.0000

TABLE 3. Results of the third experiment.

TABLE 3. Results of the third experiment.

TABLE 4

TABLE 4. Results of the fourth experiment.

TABLE 4. Results of the fourth experiment.

TABLE 4. Results of the fourth experiment.

TABLE 4. Results of the fourth experiment.

β	α	γ
0.0000	0.0000	0.0000
0.0000	0.0000	0.0000
0.0000	0.0000	0.0000
0.0000	0.0000	0.0000
0.0000	0.0000	0.0000
0.0000	0.0000	0.0000

TABLE 4. Results of the fourth experiment.

TABLE 4. Results of the fourth experiment.

Table CVIII

0.03135 N Sodium Ethylate,
0.35 N Methyl Iodide and
0.31875 N Sodium Iodide.

A = 10.00 B = 80.00

t	x	K_v
15	3.234	0.003136
20	3.880	0.003163
36	3.593	0.003193
33	4.324	0.003307
40	4.856	0.003150
49	5.648	0.003211

Average = 0.003176

$K_n = 0.06964$

Table CX

0.03135 N Sodium Ethylate,
0.25 N Methyl Iodide and
0.46875 N Sodium Iodide.

A = 9.944 B = 80.00

t	x	K_v
14	1.816	0.001811
33	2.664	0.001793
32	3.840	0.001946
40	4.544	0.001958
49	5.340	0.001971
58	5.848	0.001984

Average = 0.001911

$K_n = 0.06148$

Table CIX

0.03135 N Sodium Ethylate,
0.25 N Methyl Iodide and
0.31875 N Sodium Iodide.

A = 10.05 B = 80.00

t	x	K_v
20	2.898	0.003178
25	3.450	0.003162
30	3.963	0.003156
35	4.442	0.003159
40	4.842	0.003136
45	5.163	0.003087
50	5.594	0.003128

Average = 0.003144

$K_n = 0.06832$

Table CXI

0.03135 N Sodium Ethylate,
0.25 N Methyl Iodide and
0.46875 N Sodium Iodide.

A = 10.01 B = 80.00

t	x	K_v
15	3.010	0.001894
24	3.090	0.001965
40	4.482	0.001916
48	5.170	0.001965
56	5.594	0.001904
63	6.074	0.001939

Average = 0.001939

$K_n = 0.06199$

Table 1

Table 1 shows the results of the analysis of variance for the effect of the treatment on the yield of the different varieties of wheat. The results are given in the following table:

Treatment	Yield (kg/ha)	Standard Error
Control	2.15	0.12
Fertilizer A	2.35	0.11
Fertilizer B	2.25	0.10
Fertilizer C	2.45	0.09
Fertilizer D	2.55	0.08
Fertilizer E	2.65	0.07
Fertilizer F	2.75	0.06
Fertilizer G	2.85	0.05

Table 1 shows the results of the analysis of variance for the effect of the treatment on the yield of the different varieties of wheat.

Table 1 shows the results of the analysis of variance for the effect of the treatment on the yield of the different varieties of wheat.

Table 2

Table 2 shows the results of the analysis of variance for the effect of the treatment on the yield of the different varieties of wheat. The results are given in the following table:

Treatment	Yield (kg/ha)	Standard Error
Control	2.15	0.12
Fertilizer A	2.35	0.11
Fertilizer B	2.25	0.10
Fertilizer C	2.45	0.09
Fertilizer D	2.55	0.08
Fertilizer E	2.65	0.07
Fertilizer F	2.75	0.06
Fertilizer G	2.85	0.05

Table 2 shows the results of the analysis of variance for the effect of the treatment on the yield of the different varieties of wheat.

Table 2 shows the results of the analysis of variance for the effect of the treatment on the yield of the different varieties of wheat.

Table 3

Table 3 shows the results of the analysis of variance for the effect of the treatment on the yield of the different varieties of wheat. The results are given in the following table:

Treatment	Yield (kg/ha)	Standard Error
Control	2.15	0.12
Fertilizer A	2.35	0.11
Fertilizer B	2.25	0.10
Fertilizer C	2.45	0.09
Fertilizer D	2.55	0.08
Fertilizer E	2.65	0.07
Fertilizer F	2.75	0.06
Fertilizer G	2.85	0.05

Table 3 shows the results of the analysis of variance for the effect of the treatment on the yield of the different varieties of wheat.

Table 3 shows the results of the analysis of variance for the effect of the treatment on the yield of the different varieties of wheat.

Table 4

Table 4 shows the results of the analysis of variance for the effect of the treatment on the yield of the different varieties of wheat. The results are given in the following table:

Treatment	Yield (kg/ha)	Standard Error
Control	2.15	0.12
Fertilizer A	2.35	0.11
Fertilizer B	2.25	0.10
Fertilizer C	2.45	0.09
Fertilizer D	2.55	0.08
Fertilizer E	2.65	0.07
Fertilizer F	2.75	0.06
Fertilizer G	2.85	0.05

Table 4 shows the results of the analysis of variance for the effect of the treatment on the yield of the different varieties of wheat.

Table 4 shows the results of the analysis of variance for the effect of the treatment on the yield of the different varieties of wheat.

Table CXII

0.03135 N Sodium Ethylate,
0.35 N Methyl Iodide and
0.96875 N Sodium Iodide.

A = 9.99 B = 80.00

t	x	K_v
21	3.454	0.001704
29	3.150	0.001666
37	3.886	0.001707
46	4.486	0.001670
54	5.096	0.001710
64	5.574	0.001659

Average = 0.001686

$K_n = 0.05400$

Table CXIII

0.03125 N Sodium Ethylate,
0.35 N Methyl Iodide and
0.96875 N Sodium Iodide.

A = 10.02 B = 80.00

t	x	K_v
13	1.493	0.001562
31	3.324	0.001587
30	3.230	0.001654
38	3.788	0.001606
46	4.348	0.001597
55	5.044	0.001653
63	5.556	0.001674

Average = 0.001620

$K_n = 0.05173$

TABLE 1

PERCENTAGE OF TOTAL POPULATION

BY SEX AND AGE GROUP

1950-1959

1950-1959

AGE	MALE	FEMALE
0-4	10.1	10.1
5-9	9.2	9.2
10-14	8.3	8.3
15-19	7.4	7.4
20-24	6.5	6.5
25-29	5.6	5.6
30-34	4.7	4.7
35-39	3.8	3.8

1960-1969

1960-1969

TABLE 2

PERCENTAGE OF TOTAL POPULATION

BY SEX AND AGE GROUP

1950-1959

1950-1959

AGE	MALE	FEMALE
0-4	10.1	10.1
5-9	9.2	9.2
10-14	8.3	8.3
15-19	7.4	7.4
20-24	6.5	6.5
25-29	5.6	5.6
30-34	4.7	4.7
35-39	3.8	3.8

1960-1969

1960-1969

Table CXIV

Summary of Sodium Ethylate, Methyl Iodide and Sodium Iodide at 25°C.

V		K_n	Average K_n
Sodium ethylate	Sodium iodide		
33	33	0.08995	0.08998
		0.09003	
33	32/3	0.07795	0.07845
		0.07964	
		0.07776	
33	32/7	0.06964	0.06898
		0.06833	
33	32/15	0.06148	0.06173
		0.06199	
32	32/31	0.05400	0.05387
		0.05173	
16	16	0.08323	0.08297
		0.08387	
		0.08381	
16	16/3	0.07270	0.07276
		0.07282	
16	16/7	0.06596	0.06589
		0.06582	
16	16/15	0.05430	0.05496
		0.05561	
8	8	0.07608	0.07630
		0.07651	
8	8/3	0.06664	0.06676
		0.06689	
8	8/7	0.05765	0.05702
		0.05640	
4	4	0.06972	0.06962
		0.06952	
4	4/3	0.05915	0.05928
		0.05942	

Table CXV

Sodium Ethylate, Methyl Iodide and Sodium Iodide at 25°C.

V			α	$1 - \alpha$	K_n found	K_n calculated	Per cent difference
Sodium ethylate	Sodium iodide	Sodium ethylate in mixture					
32	32	16	0.481	0.519	0.08998	0.09191	-3.10
32	32/3	8	0.393	0.607	0.07845	0.08590	-8.67
32	32/7	4	0.312	0.688	0.06898	0.08026	-14.05
32	32/15	2	0.234	0.766	0.06173	0.07466	-17.32
32	32/31	1	0.148	0.852	0.05367	0.06886	-23.54
16	16	8	0.393	0.607	0.08297	0.08590	-3.41
16	16/3	4	0.312	0.688	0.07276	0.08026	-9.34
16	16/7	2	0.234	0.766	0.06589	0.07466	-11.74
16	16/15	1	0.148	0.852	0.05496	0.06886	-20.19
8	8	4	0.312	0.688	0.07630	0.08026	-4.93
8	8/3	2	0.234	0.766	0.06676	0.07466	-10.58
8	8/7	1	0.148	0.852	0.05702	0.06886	-17.19
4	4	2	0.234	0.766	0.06962	0.07466	- 6.75
4	4/3	1	0.148	0.852	0.05928	0.06886	-13.91

" K_n calculated" is obtained by substituting data in equation for K_n discussed on page 11.

[illegible]

TableCXVI

Sodium Ethylate, Methyl Iodide and Sodium Iodide at 25°.

V			α	$1 - \alpha$	K_n	K_n'	Per
Sodium	Sodium	Sodium	and		found	calcu-	cent.
ethylate	iodide	ethylate	α'			lated	error
		in					
		mixture					
32	32	16	0.481	0.519	0.08998	0.0902	-0.2
32	32/3	8	0.393	0.607	0.07845	0.0828	-5.2
32	32/7	4	0.312	0.688	0.06898	0.0748	-7.7
32	32/15	2	0.234	0.766	0.06173	0.0658	-6.2
32	32/31	1	0.148	0.852	0.05227	0.0548	-3.4
16	16	8	0.393	0.607	0.08297	0.0833	-0.4
16	16/3	4	0.312	0.688	0.07276	0.0752	-3.2
16	16/7	2	0.234	0.766	0.06589	0.0661	-0.3
16	16/15	1	0.148	0.852	0.05496	0.0551	-0.5
8	8	4	0.312	0.688	0.07630	0.0761	+0.3
8	8/3	2	0.234	0.766	0.06676	0.0669	-0.2
8	8/7	1	0.148	0.852	0.05702	0.0557	+2.3
4	4	2	0.234	0.766	0.06962	0.0683	+2.0
4	4/3	1	0.148	0.852	0.05928	0.0570	+4.0

" K_n' calculated" is obtained by substituting the necessary data in the equation for K_n' on page 13.

01131001

*All data subject to audit. All figures are estimates. All figures are in millions of dollars.

Year	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100
1970	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100

CONCLUSIONS.

1. It has been shown that methyl iodide reacts at 0° with sodium, potassium and lithium ethylates. The velocity can be expressed as functions of the concentrations of both the ethylate ions and the nonionized salts.
2. If the ethylate ion reacts independently with the methyl iodide, we should obtain from all three ethylates the same value for K_1 , the reaction velocity of unit concentrations of the ethylate ions and methyl iodide. But different values might be found for the reaction velocities of the three nonionized ethylates, which, of course, do not have the same physical and chemical properties. In confirmation of this theory, we find the values 0.00535, 0.00486 and 0.00533 for K_1 , and the values 0.00337, 0.00303 and 0.00199 for K_m .
3. The "salt effect" produced by sodium iodide on the reaction of sodium ethylate and methyl iodide at 35° , has been studied in various concentrations. It has been shown that, besides the "normal salt effect" due to the change in the ionization of the sodium ethylate, produced by the added sodium iodide, there is in addition, a slight "negative abnormal salt effect" which can be expressed as a function of the concentrations of the ethylate ions and of the iodide ions.

BIOGRAPHY.

Bessie Marion Brown, the author of this dissertation, was born in Rye, New Hampshire, July 31, 1884. She received her early education in the public schools of that town, her high school training at Newburyport, Massachusetts, and the degree of B.S. from Simmons College, Boston, Massachusetts, in 1907. After holding an instructorship in chemistry at the latter institution for four years, she entered Johns Hopkins University in 1911 as a graduate student in chemistry with physical chemistry and elementary applied electricity as minor subjects. In 1911-13 she was appointed a university scholar.





